

中華民國輸入植物或植物產品檢疫規定部分規定修正對照表

修正規定				現行規定				說明
乙、有條件輸入植物或植物產品之檢疫條件 依據：植物防疫檢疫法第十四條第一項第二款、第三款規定。 一、植物或植物產品之名稱及其部位、國家或地區及其輸入檢疫條件如下表：				乙、有條件輸入植物或植物產品之檢疫條件 依據：植物防疫檢疫法第十四條第一項第二款、第三款規定。 一、植物或植物產品之名稱及其部位、國家或地區及其輸入檢疫條件如下表：				配合增訂之乙、十七點「繁殖用生植株」之定義，修正現行檢疫條件欄2之「種植用種子」為「繁殖用種子」。
植物名稱及其部位	國家或地區	病蟲害名稱	檢疫條件	植物名稱及其部位	國家或地區	病蟲害名稱	檢疫條件	
23. 下列植物生植株之全部、部分： (1) 冬瓜 (Benincasa hispida) (2) 西瓜 (Citrullus lanatus) (3) 甜瓜 (Cucumis melo) (4) 胡瓜 (Cucumis sativus) (5) 南瓜 (Cucurbita moschata) (6) 美國南瓜 (Cucurbita pepo)	亞洲及太平洋地區 (1) 澳大利亞 (昆士蘭州) (2) 中國大陸 (3) 關島 (4) 以色列 (5) 日本 (6) 南韓 (7) 北馬里亞納群島 (8) 泰國 (9) 土耳其 北美 (10) 美國(阿拉巴馬州、阿肯色州、加利福尼亞州、德拉瓦州、佛羅里達州、喬治亞州、伊利諾州、印第安那州、愛荷華州、馬里蘭州、密西西比州、密蘇里州、北卡羅來那州、奧克拉荷馬州、奧勒岡州、南卡羅來那州、德克薩斯州) 中南美 (11) 巴西	細菌性果斑病 (Acidovorax avenae subsp. citrulli (Schaad et al.) Willemet al. (Pseudomonas pseudoalcaligenes subsp. citrulli Schaad et al.))	1、非繁殖性之植物部分、果實及供食用之種子應檢附輸出國植物檢疫機關(構)簽發之植物檢疫證明書，證明未染細菌性果斑病。否則銷燬或退運。 2、植株及繁殖用種子應檢附輸出國植物檢疫機關(構)簽發之植物檢疫證明書，證明經田間檢疫未染細菌性果斑病，或於輸出前經實驗室檢測未染細菌性果斑病，或於輸出前經適當之檢疫處理。	23. 下列植物生植株之全部、部分： (1) 冬瓜 (Benincasa hispida) (2) 西瓜 (Citrullus lanatus) (3) 甜瓜 (Cucumis melo) (4) 胡瓜 (Cucumis sativus) (5) 南瓜 (Cucurbita moschata) (6) 美國南瓜 (Cucurbita pepo)	亞洲及太平洋地區 (4) 澳大利亞 (昆士蘭州) (5) 中國大陸 (6) 關島 (4) 以色列 (5) 日本 (6) 南韓 (7) 北馬里亞納群島 (8) 泰國 (9) 土耳其 北美 (10) 美國(阿拉巴馬州、阿肯色州、加利福尼亞州、德拉瓦州、佛羅里達州、喬治亞州、伊利諾州、印第安那州、愛荷華州、馬里蘭州、密西西比州、密蘇里州、北卡羅來那州、奧克拉荷馬州、奧勒岡州、南卡羅來那州、德克薩斯州) 中南美 (11) 巴西	細菌性果斑病 (Acidovorax avenae subsp. citrulli (Schaad et al.) Willemet al. (Pseudomonas pseudoalcaligenes subsp. citrulli Schaad et al.))	1、非繁殖性之植物部分、果實及供食用之種子應檢附輸出國植物檢疫機關(構)簽發之植物檢疫證明書，證明未染細菌性果斑病。否則銷燬或退運。 2、植株及種植用種子應檢附輸出國植物檢疫機關(構)簽發之植物檢疫證明書，證明經田間檢疫未染細菌性果斑病，或於輸出前經實驗室檢測未染細菌性果斑病，或於輸出前經適當之檢疫處理。	
植物名稱及其部位	國家或地區	病蟲害名稱	檢疫條件	植物名稱及其部位	國家或地區	病蟲害名稱	檢疫條件	一、配合增訂之乙、十七點「繁殖用生植株」之定義，「繁殖用地上部」已納入「繁殖用生植株範圍」，另切花及果實非屬
37. 植物名稱及其部位 I. 下列繁殖用生植	亞洲及太平洋地區 (1) 阿富汗	梨花枯病 (Pseudomonas	1. 應檢附輸出國植物檢疫機關(構)	37. 植物名稱及其部位 I. 下列繁殖用之生	亞洲及太平洋地區 (1) 阿富汗	梨花枯病 (Pseudomona	1. 應檢附輸出國植物檢疫機關(構)	

<p>株 (種子除外) :</p> <p>(1) 黃秋葵 (<i>Abelmoschus esculentus = Hibiscus esculentus</i>)</p> <p>(2) 烏頭屬 (<i>Aconitum</i> spp.)</p> <p>(3) 臺灣羊桃 (<i>Actinidia chinensis</i>)</p> <p>(4) 奇異果 (<i>Actinidia deliciosa</i>)</p> <p>(5) 赤楊屬之一種 (<i>Alnus glutinosa</i>)</p> <p>(6) 梔木屬 (<i>Aralia</i> spp.)</p> <p>(7) 漿果鵝屬 (<i>Arbutus</i> spp.)</p> <p>(8) 歐洲白樺 (<i>Betula pendula</i>)</p> <p>(9) 旋花科之一種 (<i>Calystegia sepium</i>)</p> <p>(10) 番椒 (<i>Capsicum annuum</i>)</p> <p>(11) 紅花 (<i>Carthamus tinctorius</i>)</p> <p>(12) 矢車菊屬之一種 (<i>Centaurea stoebe</i>)</p> <p>(13) 野菊 (<i>Chrysanthemum</i>)</p>	<p>(2) 亞塞拜然</p> <p>(3) 澳大利亞</p> <p>(4) 孟加拉</p> <p>(5) 白俄羅斯</p> <p>(6) 中國大陸</p> <p>(7) 喬治亞</p> <p>(8) 印度</p> <p>(9) 伊朗</p> <p>(10) 以色列</p> <p>(11) 日本</p> <p>(12) 北韓</p> <p>(13) 南韓</p> <p>(14) 吉爾吉斯</p> <p>(15) 黎巴嫩</p> <p>(16) 紐西蘭</p> <p>(17) 俄羅斯聯邦</p> <p>(18) 巴基斯坦</p> <p>(19) 斯里蘭卡</p> <p>(20) 泰國</p> <p>(21) 土耳其</p> <p>(22) 烏茲別克 非洲</p> <p>(23) 阿爾及利亞</p> <p>(24) 埃及</p> <p>(25) 衣索比亞</p> <p>(26) 肯亞</p> <p>(27) 賴索托</p> <p>(28) 利比亞</p> <p>(29) 馬拉威</p> <p>(30) 摩洛哥</p> <p>(31) 奈及利亞</p> <p>(32) 南非</p> <p>(33) 坦尚尼亞</p> <p>(34) 突尼西亞</p> <p>(35) 烏干達</p> <p>(36) 辛巴威 歐洲</p> <p>(37) 奧地利</p> <p>(38) 比利時</p> <p>(39) 波士尼亞赫</p>	<p><i>syringae</i> pv. <i>syringae</i>)</p> <p>簽發之植物檢疫證明書，證明經田間檢疫未罹染 <i>Pseudomonas syringae</i> pv. <i>syringae</i>，或於輸出前經實驗室檢測未罹染 <i>Pseudomonas syringae</i> pv. <i>syringae</i>，否則銷燬或退運。</p> <p>2. 日本產梨接穗依「日本產梨接穗輸入檢疫條件」辦理輸入。</p> <p>3. 中國大陸產梨接穗依「中國大陸產梨接穗輸入檢疫條件」辦理輸入。</p>	<p>植株及地上部(切花、種子及果實除外) :</p> <p>(1) 黃秋葵 (<i>Abelmoschus esculentus = Hibiscus esculentus</i>)</p> <p>(2) 烏頭屬 (<i>Aconitum</i> spp.)</p> <p>(3) 臺灣羊桃 (<i>Actinidia chinensis</i>)</p> <p>(4) 奇異果 (<i>Actinidia deliciosa</i>)</p> <p>(5) 赤楊屬之一種 (<i>Alnus glutinosa</i>)</p> <p>(6) 梔木屬 (<i>Aralia</i> spp.)</p> <p>(7) 漿果鵝屬 (<i>Arbutus</i> spp.)</p> <p>(8) 歐洲白樺 (<i>Betula pendula</i>)</p> <p>(9) 旋花科之一種 (<i>Calystegia sepium</i>)</p> <p>(10) 番椒 (<i>Capsicum annuum</i>)</p> <p>(11) 紅花 (<i>Carthamus tinctorius</i>)</p> <p>(12) 矢車菊屬之一種 (<i>Centaurea stoebe</i>)</p>	<p>(2) 亞塞拜然</p> <p>(3) 澳大利亞</p> <p>(4) 孟加拉</p> <p>(5) 白俄羅斯</p> <p>(6) 中國大陸</p> <p>(7) 喬治亞</p> <p>(8) 印度</p> <p>(9) 伊朗</p> <p>(10) 以色列</p> <p>(11) 日本</p> <p>(12) 北韓</p> <p>(13) 南韓</p> <p>(14) 吉爾吉斯</p> <p>(15) 黎巴嫩</p> <p>(16) 紐西蘭</p> <p>(17) 俄羅斯聯邦</p> <p>(18) 巴基斯坦</p> <p>(19) 斯里蘭卡</p> <p>(20) 泰國</p> <p>(21) 土耳其</p> <p>(22) 烏茲別克 非洲</p> <p>(23) 阿爾及利亞</p> <p>(24) 埃及</p> <p>(25) 衣索比亞</p> <p>(26) 肯亞</p> <p>(27) 賴索托</p> <p>(28) 利比亞</p> <p>(29) 馬拉威</p> <p>(30) 摩洛哥</p> <p>(31) 奈及利亞</p> <p>(32) 南非</p> <p>(33) 坦尚尼亞</p> <p>(34) 突尼西亞</p> <p>(35) 烏干達</p> <p>(36) 辛巴威 歐洲</p> <p>(37) 奧地利</p> <p>(38) 比利時</p> <p>(39) 波士尼亞赫塞</p>	<p><i>s. syringae</i> pv. <i>syringae</i>)</p> <p>簽發之植物檢疫證明書，證明經田間檢疫未罹染 <i>Pseudomonas syringae</i> pv. <i>syringae</i>，或於輸出前經實驗室檢測未罹染 <i>Pseudomonas syringae</i> pv. <i>syringae</i>，否則銷燬或退運。</p> <p>2. 日本產梨接穗依「日本產梨接穗輸入檢疫條件」辦理輸入。</p> <p>3. 中國大陸產梨接穗依「中國大陸產梨接穗輸入檢疫條件」辦理輸入。</p>	<p>繁殖用生植株，爰修正植物名稱及其部位欄 I 之序文文字，修正前後本項規範之植物部位範圍並未改變。</p> <p>二、另統一法規用詞，修正植物名稱及其部位欄 II 之序文種植用種子為繁殖用種子。</p>
--	--	--	--	---	---	---

<i>indicum</i>)	塞哥維納			(13)野菊	哥維納		
(14)樟樹	(40) 保加利亞			(<i>Chrysanthemum</i>	(40) 保加利亞		
(<i>Cinnamomum</i>	(41) 賽普勒斯			<i>indicum</i>)	(41) 賽普勒斯		
<i>camphora</i>)	(42) 捷克			(14)樟樹	(42) 捷克		
(15)柑橘屬	(43) 丹麥			(<i>Cinnamomum</i>	(43) 丹麥		
(<i>Citrus spp.</i>)	(44) 法國			<i>camphora</i>)	(44) 法國		
(16)阿拉比卡咖啡	(45) 德國			(15)柑橘屬	(45) 德國		
(<i>Coffea arabica</i>	(46) 希臘			(<i>Citrus spp.</i>)	(46) 希臘		
)	(47) 匈牙利			(16)阿拉比卡咖啡	(47) 匈牙利		
(17)胡瓜	(48) 愛爾蘭			(<i>Coffea arabica</i>	(48) 愛爾蘭		
(<i>Cucumis</i>	(49) 義大利)	(49) 義大利		
<i>sativus</i>)	(50) 拉脫維亞			(17)胡瓜	(50) 拉脫維亞		
(18)南瓜屬	(51) 立陶宛			(<i>Cucumis</i>	(51) 立陶宛		
(<i>Cucurbita spp.</i>)	(52) 馬其頓			<i>sativus</i>)	(52) 馬其頓		
(19)瓜爾豆	(53) 摩爾多瓦			(18)南瓜屬	(53) 摩爾多瓦		
(<i>Cyamopsis te-t</i>	(54) 荷蘭			(<i>Cucurbita spp.</i>)	(54) 荷蘭		
<i>ragonoloba</i>)	(55) 波蘭			(19)瓜爾豆	(55) 波蘭		
(20)榲桲	(56) 葡萄牙			(<i>Cyamopsis te-t</i>	(56) 葡萄牙		
(<i>Cydonia</i>	(57) 羅馬尼亞			<i>ragonoloba</i>)	(57) 羅馬尼亞		
<i>oblonga</i>)	(58) 塞爾維亞			(20)榲桲	(58) 塞爾維亞		
(21)大理花	(59) 斯洛維尼亞			(<i>Cydonia</i>	(59) 斯洛維尼亞		
(<i>Dahlia</i>	(60) 西班牙			<i>oblonga</i>)	(60) 西班牙		
<i>pinnata</i>)	(61) 瑞典			(21)大理花	(61) 瑞典		
(22)瑞香	(62) 瑞士			(<i>Dahlia</i>	(62) 瑞士		
(<i>Daphne</i>	(63) 英國			<i>pinnata</i>)	(63) 英國		
<i>odora</i>)	(64) 烏克蘭			(22)瑞香	(64) 烏克蘭		
(23)柿	北美			(<i>Daphne</i>	北美		
(<i>Diospyros</i>	(65) 加拿大			<i>odora</i>)	(65) 加拿大		
<i>kaki</i>)	(66) 墨西哥			(23)柿	(66) 墨西哥		
(24)連翹屬之一	(67) 美國			(<i>Diospyros</i>	(67) 美國		
種	中南美			<i>kaki</i>)	中南美		
(<i>Forsythia</i>	(68) 阿根廷			(24)連翹屬之一	(68) 阿根廷		
<i>intermedia</i>)	(69) 巴貝多			種	(69) 巴貝多		
(25)金柑屬	(70) 巴西			(<i>Forsythia</i>	(70) 巴西		
(<i>Fortunella</i>	(71) 智利			<i>intermedia</i>)	(71) 智利		
<i>spp.</i>)	(72) 薩爾瓦多			(25)金柑屬	(72) 薩爾瓦多		
(26)歐洲草莓	(73) 瓜地馬拉			(<i>Fortunella</i>	(73) 瓜地馬拉		
(<i>Fragaria</i>	(74) 宏都拉斯			<i>spp.</i>)	(74) 宏都拉斯		
<i>vesca</i>)	(75) 巴拿馬			(26)歐洲草莓	(75) 巴拿馬		
	(76) 波多黎各				(76) 波多黎各		

(27)大豆 (<i>Glycine max</i>)	(77) 烏拉圭 (78) 委內瑞拉			(<i>Fragaria vesca</i>)	(77) 烏拉圭 (78) 委內瑞拉		
(28)木槿屬 (<i>Hibiscus spp.</i>)				(27)大豆 (<i>Glycine max</i>)			
(29)大麥 (<i>Hordeum vulgare</i>)				(28)木槿屬 (<i>Hibiscus spp.</i>)			
(30)胡桃 (<i>Juglans regia</i>)				(29)大麥 (<i>Hordeum vulgare</i>)			
(31)白花鵲豆 (<i>Lablab purpureus</i>)				(30)胡桃 (<i>Juglans regia</i>)			
(32)女貞屬之一種 (<i>Ligustrum ovalifolium</i>)				(31)白花鵲豆 (<i>Lablab purpureus</i>)			
(33)荔枝 (<i>Litchi chinensis</i>)				(32)女貞屬之一種 (<i>Ligustrum ovalifolium</i>)			
(34)木蘭屬 (<i>Magnolia spp.</i>)				(33)荔枝 (<i>Litchi chinensis</i>)			
(35)檬果 (<i>Mangifera indica</i>)				(34)木蘭屬 (<i>Magnolia spp.</i>)			
(36)蘋果 (<i>Malus domestica</i>)				(35)檬果 (<i>Mangifera indica</i>)			
(37)虎爪豆 (<i>Mucuna pruriens</i>)				(36)蘋果 (<i>Malus domestica</i>)			
(38)芭蕉屬之一種 (<i>Musa paradisiaca</i>)				(37)虎爪豆 (<i>Mucuna pruriens</i>)			
(39)歐洲夾竹桃 (<i>Nerium oleander</i>)				(38)芭蕉屬之一種 (<i>Musa paradisiaca</i>)			
(40)菸草 (<i>Nicotiana tabacum</i>)				(39)歐洲夾竹桃 (<i>Nerium oleander</i>)			
				(40)菸草			

(41) 橄欖 (<i>Olea europaea</i>)				(<i>Nicotiana tabacum</i>)				
(42) 稻 (<i>Oryza sativa</i>)				(41) 橄欖 (<i>Olea europaea</i>)				
(43) 百香果 (<i>Passiflora edulis</i>)				(42) 稻 (<i>Oryza sativa</i>)				
(44) 酪梨 (<i>Persea americana</i>)				(43) 百香果 (<i>Passiflora edulis</i>)				
(45) 花豆 (<i>Phaseolus coccineus</i>)				(44) 酪梨 (<i>Persea americana</i>)				
(46) 菜豆 (<i>Phaseolus lunatus</i>)				(45) 花豆 (<i>Phaseolus coccineus</i>)				
(47) 敏豆 (<i>Phaseolus vulgaris</i>)				(46) 菜豆 (<i>Phaseolus lunatus</i>)				
(48) 山梅花屬 (<i>Philadelphus</i> spp.)				(47) 敏豆 (<i>Phaseolus vulgaris</i>)				
(49) 放射松 (<i>Pinus radiata</i>)				(48) 山梅花屬 (<i>Philadelphus</i> spp.)				
(50) 胡椒 (<i>Piper nigrum</i>)				(49) 放射松 (<i>Pinus radiata</i>)				
(51) 豌豆 (<i>Pisum sativum</i>)				(50) 胡椒 (<i>Piper nigrum</i>)				
(52) 楊屬 (<i>Populus</i> spp.)				(51) 豌豆 (<i>Pisum sativum</i>)				
(53) 李屬 (<i>Prunus</i> spp.)				(52) 楊屬 (<i>Populus</i> spp.)				
(54) 梨屬 (<i>Pyrus</i> spp.)				(53) 李屬 (<i>Prunus</i> spp.)				
(55) 杜鵑花屬 (<i>Rhododendron</i> spp.)				(54) 梨屬 (<i>Pyrus</i> spp.)				
(56) 薔薇屬 (<i>Rosa</i> spp.)				(55) 杜鵑花屬 (<i>Rhododendron</i> spp.)				

<p>(57) 柳屬 (<i>Salix</i> spp.)</p> <p>(58) 樹番茄 (<i>Solanum</i> <i>betaceum</i> = <i>Cyphomandra</i> <i>betacea</i>)</p> <p>(59) 番茄 (<i>Solanum</i> <i>lycopersicum</i> =<i>Lycopersium</i> <i>esculentum</i>)</p> <p>(60) 高粱 (<i>Sorghum</i> <i>bicolor</i>)</p> <p>(61) 小麥 (<i>Triticum</i> <i>aestivum</i>)</p> <p>(62) 油桐 (<i>Vernicia</i> <i>fordii</i>=<i>Aleurites</i> <i>fordii</i>)</p> <p>(63) 蠶豆屬 (<i>Vicia</i> spp.)</p> <p>(64) 豇豆屬 (<i>Vigna</i> spp.)</p> <p>(65) 葡萄屬 (<i>Vitis</i> spp.)</p> <p>(66) 玉米 (<i>Zea mays</i>)</p> <p>II. 下列<u>繁</u>植用種子： (1) 瓜爾豆 (<i>Cyamopsis tetragonoloba</i>) (2) 大豆 (<i>Glycine max</i>) (3) 白花鵲豆 (<i>Lablab purpureus</i>)</p>					<p>(56) 薔薇屬 (<i>Rosa</i> spp.)</p> <p>(57) 柳屬 (<i>Salix</i> spp.)</p> <p>(58) 樹番茄 (<i>Solanum</i> <i>betaceum</i> = <i>Cyphomandra</i> <i>betacea</i>)</p> <p>(59) 番茄 (<i>Solanum</i> <i>lycopersicum</i> =<i>Lycopersium</i> <i>esculentum</i>)</p> <p>(60) 高粱 (<i>Sorghum</i> <i>bicolor</i>)</p> <p>(61) 小麥 (<i>Triticum</i> <i>aestivum</i>)</p> <p>(62) 油桐 (<i>Vernicia</i> <i>fordii</i>=<i>Aleurites</i> <i>fordii</i>)</p> <p>(63) 蠶豆屬 (<i>Vicia</i> spp.)</p> <p>(64) 豇豆屬 (<i>Vigna</i> spp.)</p> <p>(65) 葡萄屬 (<i>Vitis</i> spp.)</p> <p>(66) 玉米 (<i>Zea mays</i>)</p> <p>II. 下列<u>種</u>植用種子： (12) 瓜爾豆 (<i>Cyamopsis tetragonoloba</i>) (13) 大豆 (<i>Glycine max</i>) (14) 白花鵲豆</p>				
---	--	--	--	--	---	--	--	--	--

<p>(4) 稻 (<i>Oryza sativa</i>)</p> <p>(5) 花豆 (<i>Phaseolus coccineus</i>)</p> <p>(6) 菜豆 (<i>Phaseolus lunatus</i>)</p> <p>(7) 敏豆 (<i>Phaseolus vulgaris</i>)</p> <p>(8) 高粱 (<i>Sorghum bicolor</i>)</p> <p>(9) 小麥 (<i>Triticum aestivum</i>)</p> <p>(10) 蠶豆屬 (<i>Vicia spp.</i>)</p> <p>(11) 豇豆屬 (<i>Vigna spp.</i>)</p>				<p>(<i>Lablab purpureus</i>)</p> <p>(15) 稻 (<i>Oryza sativa</i>)</p> <p>(16) 花豆 (<i>Phaseolus coccineus</i>)</p> <p>(17) 菜豆 (<i>Phaseolus lunatus</i>)</p> <p>(18) 敏豆 (<i>Phaseolus vulgaris</i>)</p> <p>(19) 高粱 (<i>Sorghum bicolor</i>)</p> <p>(20) 小麥 (<i>Triticum aestivum</i>)</p> <p>(21) 蠶豆屬 (<i>Vicia spp.</i>)</p> <p>(22) 豇豆屬 (<i>Vigna spp.</i>)</p>				
<p>39.下列繁殖用生植株 <u>(種子除外)</u>：</p> <p>(1) 西洋芹 (<i>Apium graveolens</i>)</p> <p>(2) 辣根 (<i>Armoracia rusticana</i>)</p> <p>(3) 甜菜 (<i>Beta vulgaris</i>)</p> <p>(4) 胡瓜 (<i>Cucumis sativus</i>)</p> <p>(5) 豆科 (<i>Fabaceae</i>)</p> <p>(6) 連翹 (<i>Forsythia suspense</i>)</p>	<p>亞洲及太平洋地區</p> <p>(1) 澳大利亞</p> <p>(2) 紐西蘭</p> <p>(3) 俄羅斯聯邦</p> <p>(4) 土耳其</p> <p>(5) 烏克蘭</p> <p>非洲</p> <p>(6) 南非</p> <p>歐洲</p> <p>(7) 比利時</p> <p>(8) 保加利亞</p> <p>(9) 捷克</p> <p>(10) 丹麥</p> <p>(11) 法國</p> <p>(12) 德國</p> <p>(13) 挪威</p>	<p>南芥嵌紋病毒 (<i>Arabis mosaic virus</i> (Nepovirus))</p>	<p>應檢附輸出國植物檢疫機關(構)簽發之植物檢疫證明書，證明經田間檢疫或於輸出前經實驗室檢測未罹染南芥嵌紋病毒，否則銷燬或退運。</p>	<p>39.下列繁殖用生植株：</p> <p>(1) 西洋芹 (<i>Apium graveolens</i>)</p> <p>(2) 山葵 (<i>Armoracia rusticana</i>)</p> <p>(3) 甜菜 (<i>Beta vulgaris</i>)</p> <p>(4) 胡瓜 (<i>Cucumis sativus</i>)</p> <p>(5) 豆科 (<i>Fabaceae</i>)</p> <p>(6) 連翹 (<i>Forsythia suspense</i>)</p>	<p>亞洲及太平洋地區</p> <p>(1) 澳大利亞</p> <p>(2) 紐西蘭</p> <p>(3) 俄羅斯聯邦</p> <p>(4) 土耳其</p> <p>(5) 烏克蘭</p> <p>非洲</p> <p>(6) 南非</p> <p>歐洲</p> <p>(7) 比利時</p> <p>(8) 保加利亞</p> <p>(9) 捷克</p> <p>(10) 丹麥</p> <p>(11) 法國</p> <p>(12) 德國</p> <p>(13) 挪威</p>	<p>南芥嵌紋病毒 (<i>Arabis mosaic virus</i> (Nepovirus))</p>	<p>應檢附輸出國植物檢疫機關(構)簽發之植物檢疫證明書，證明經田間檢疫或於輸出前經實驗室檢測未罹染南芥嵌紋病毒，否則銷燬或退運。</p>	<p>一、原規範之「繁殖用生植株」即未包含種子，配合增訂之乙、十七點「繁殖用生植株」之定義，爰酌修植物部位部分文字，修正前後本項規範之植物部位範圍並未改變。</p> <p>二、植物名稱及其部位欄(2)山葵，原中文名有誤，修正名稱為辣根。</p>

(7) 草莓 (<i>Fragaria × ananassa</i>)	(14) 匈牙利			(7) 草莓 (<i>Fragaria × ananassa</i>)	(14) 匈牙利		
(8) 唐菖蒲 (<i>Gladiolus hybridus</i>)	(15) 愛爾蘭			(8) 唐菖蒲 (<i>Gladiolus hybridus</i>)	(15) 愛爾蘭		
(9) 蛇麻 (<i>Humulus lupulus</i>)	(16) 義大利			(9) 蛇麻 (<i>Humulus lupulus</i>)	(16) 義大利		
(10) 萵苣 (<i>Lactuca sativa</i>)	(17) 盧森堡			(10) 萵苣 (<i>Lactuca sativa</i>)	(17) 盧森堡		
(11) 百合屬 (<i>Lilium spp.</i>)	(18) 摩爾達維亞			(11) 百合屬 (<i>Lilium spp.</i>)	(18) 摩爾達維亞		
(12) 橄欖 (<i>Olea europaea</i>)	(19) 荷蘭			(12) 橄欖 (<i>Olea europaea</i>)	(19) 荷蘭		
(13) 水仙屬 (<i>Narcissus spp.</i>)	(20) 波蘭			(13) 水仙屬 (<i>Narcissus spp.</i>)	(20) 波蘭		
(14) 甜櫻桃 (<i>Prunus avium</i>)	(21) 羅馬尼亞			(14) 甜櫻桃 (<i>Prunus avium</i>)	(21) 羅馬尼亞		
(15) 杏 (<i>Prunus armeniaca</i>)	(22) 南斯拉夫			(15) 杏 (<i>Prunus armeniaca</i>)	(22) 南斯拉夫		
(16) 歐洲酸櫻桃 (<i>Prunus cerasus</i>)	(23) 斯洛維尼亞			(16) 歐洲酸櫻桃 (<i>Prunus cerasus</i>)	(23) 斯洛維尼亞		
(17) 歐洲李 (<i>Prunus domestica</i>)	亞			(17) 歐洲李 (<i>Prunus domestica</i>)	(24) 西班牙		
(18) 扁桃 (<i>Prunus dulcis</i>)	(24) 西班牙			(18) 扁桃 (<i>Prunus dulcis</i>)	(25) 瑞典		
(19) 桂櫻 (<i>Prunus laurocerasus</i>)	(25) 瑞典			(19) 桂櫻 (<i>Prunus laurocerasus</i>)	(26) 瑞士		
(20) 桃 (<i>Prunus persica</i>)	(26) 瑞士			(20) 桃 (<i>Prunus persica</i>)	(27) 芬蘭		
(21) 大黃 (<i>Rheum palmatum</i>)	(27) 芬蘭			(21) 大黃 (<i>Rheum palmatum</i>)	(28) 英國		
(22) 薔薇屬 (<i>Rosa spp.</i>)	(28) 英國			(22) 薔薇屬 (<i>Rosa spp.</i>)	(29) 烏克蘭		
(23) 覆盆子 (<i>Rubus idaeus</i>)	(29) 烏克蘭			(23) 覆盆子 (<i>Rubus idaeus</i>)	北美		
(24) 西洋接骨木	北美			(24) 西洋接骨木	(30) 加拿大		
	(30) 加拿大						

(<i>Sambucus nigra</i>) (25) 白花苜蓿 (<i>Trifolium repens</i>) (26) 葡萄 (<i>Vitis vinifera</i>)				(<i>Sambucus nigra</i>) (25) 白花苜蓿 (<i>Trifolium repens</i>) (26) 葡萄 (<i>Vitis vinifera</i>)				
40.植物名稱及其部位： I.下列繁殖用生植株 (種子除外)： (1)曼陀羅木屬 (<i>Brugmansia</i> spp.) (2)夜香木屬 (<i>Cestrum</i> spp.) (3)大理花屬 (<i>Dahlia</i> spp.) (4)曼陀羅屬 (<i>Datura</i> spp.) (5)藍花茄 (<i>Lycianthes rantonnetii</i>) (6)酪梨 (<i>Persea americana</i>) (7)好望角醋栗 (<i>Physalis peruviana</i>) (8)果醬木 (<i>Streptosolen jamesonii</i>) II.下列繁殖用生植株： (1)番椒屬 (<i>Capsicum</i> spp.) (2)矮牽牛屬 (<i>Petunia</i> spp.) (3)茄屬 (<i>Solanum</i> spp.)	亞洲及太平洋地區 (1) 阿富汗 (2) 澳大利亞 (3) 亞塞拜然 (4) 孟加拉 (5) 中國大陸 (6) 喬治亞 (7) 印度 (8) 伊朗 (9) 以色列 (10) 日本 (11) 土耳其 (12) 哈薩克 (13) 巴基斯坦 非洲 (14) 埃及 (15) 奈及利亞 (16) 迦納 (17) 烏干達 (18) 肯亞 歐洲 (19) 奧地利 (20) 白俄羅斯 (21) 比利時 (22) 克羅埃西亞 (23) 捷克 (24) 德國 (25) 希臘 (26) 匈牙利 (27) 義大利 (28) 馬爾他 (29) 蒙特內哥羅 (30) 荷蘭 (31) 波蘭	馬鈴薯紡錘形塊莖類病毒 (<i>Potato spindle tuber viroid</i> , PSTVd)	應檢附輸出國植物檢疫機關(構)簽發之植物檢疫證明書，證明於輸出前經實驗室檢測未罹染馬鈴薯紡錘形塊莖類病毒，否則銷燬或退運。	40.植物名稱及其部位： I.下列繁殖用生植株： (1)曼陀羅木屬 (<i>Brugmansia</i> spp.) (2)夜香木屬 (<i>Cestrum</i> spp.) (3)大理花屬 (<i>Dahlia</i> spp.) (4)曼陀羅屬 (<i>Datura</i> spp.) (5)藍花茄 (<i>Lycianthes rantonnetii</i>) (6)酪梨 (<i>Persea americana</i>) (7)好望角醋栗 (<i>Physalis peruviana</i>) (8)果醬木 (<i>Streptosolen jamesonii</i>) II.下列繁殖用生植株及種子： (1)番椒屬 (<i>Capsicum</i> spp.) (2)矮牽牛屬 (<i>Petunia</i> spp.) (3)茄屬 (<i>Solanum</i> spp.)	亞洲及太平洋地區 (1) 阿富汗 (2) 澳大利亞 (3) 亞塞拜然 (4) 孟加拉 (5) 中國大陸 (6) 喬治亞 (7) 印度 (8) 伊朗 (9) 以色列 (10) 日本 (11) 紐西蘭 非洲 (12) 土耳其 (13) 埃及 (14) 奈及利亞 歐洲 (15) 奧地利 (16) 白俄羅斯 (17) 比利時 (18) 克羅埃西亞 (19) 捷克 (20) 德國 (21) 希臘 (22) 匈牙利 (23) 義大利 (24) 馬爾他 (25) 蒙特內哥羅 (26) 荷蘭 (27) 波蘭 (28) 俄羅斯聯邦 (29) 斯洛維尼亞 (30) 西班牙 (31) 英國 (32) 烏克蘭	馬鈴薯紡錘形塊莖類病毒 (<i>Potato spindle tuber viroid</i> , PSTVd)	應檢附輸出國植物檢疫機關(構)簽發之植物檢疫證明書，證明於輸出前經實驗室檢測未罹染馬鈴薯紡錘形塊莖類病毒，否則銷燬或退運。	一、查本項規定 I.原規範之「繁殖用生植株」即未包含種子，II.原規範之「繁殖用生植株及種子」之植物部位包含種子，配合增訂之乙、十七點「繁殖用生植株」之定義，爰酌修文字，修正前後本項規範之植物部位範圍並未改變。 二、配合刪除乙、第十六點及其附件「茄屬及番椒屬種子輸入檢疫條件」規定，並依查詢科學文獻結果，修正國家或地區欄，增列五個發生國家，並刪除紐西蘭，另序號配合調整。

	<p>(32) 俄羅斯聯邦</p> <p>(33) 斯洛維尼亞</p> <p>(34) 西班牙</p> <p>(35) 英國</p> <p>(36) 烏克蘭</p> <p>北美</p> <p>(37) 墨西哥</p> <p>中南美</p> <p>(38) 巴西</p> <p>(39) 哥斯大黎加</p> <p>(40) 多明尼加</p> <p>(41) 秘魯</p> <p>(42) 委內瑞拉</p>				<p>北美</p> <p>(33) 墨西哥</p> <p>中南美</p> <p>(34) 巴西</p> <p>(35) 哥斯大黎加</p> <p>(36) 多明尼加</p> <p>(37) 秘魯</p> <p>(38) 委內瑞拉</p>			
<p>41.下列繁殖用生植株：</p> <p>(1) 曼陀羅 (<i>Datura stramonium</i>)</p> <p>(2) 羅勒 (<i>Ocimum basilicum</i>)</p> <p>(3) 茄屬 (<i>Solanum</i> spp.)</p>	<p>亞洲及太平洋地區</p> <p>(1) 敘利亞</p> <p>(2) 土耳其</p> <p>(3) 中國大陸</p> <p>(4) 以色列</p> <p>(5) 土耳其</p> <p>非洲</p> <p>(6) 卡那利群島 (西班牙屬)</p> <p>(7) 南非</p> <p>(8) 埃及</p> <p>(9) 摩洛哥</p> <p>歐洲</p> <p>(10) 奧地利</p> <p>(11) 比利時</p> <p>(12) 保加利亞</p> <p>(13) 賽普勒斯</p> <p>(14) 丹麥</p> <p>(15) 法國</p> <p>(16) 德國</p> <p>(17) 希臘</p> <p>(18) 匈牙利</p> <p>(19) 愛爾蘭</p> <p>(20) 義大利</p> <p>(21) 立陶宛</p> <p>(22) 荷蘭</p>	<p>香瓜茄嵌紋病毒(<i>Pepino mosaic virus</i> (<i>Potexvirus</i>))</p>	<p>1、繁殖用生植株應檢附輸出國植物檢疫機關(構)簽發之植物檢疫證明書,證明經田間檢疫或於輸出前經實驗室檢測未染香瓜茄嵌紋病毒,否則銷燬或退運。</p> <p>2、繁殖用種子應檢附輸出國植物檢疫機關(構)簽發之植物檢疫證明書,證明下列條件之一,否則銷燬或退運：</p> <p>(1)經田間檢疫未染香瓜茄嵌紋病毒,並於採種後經磷酸三鈉檢疫處理。</p> <p>(2)輸出前經實驗室檢測未染香瓜茄嵌紋病毒。</p>	<p>41.下列繁殖用生植株及種子：</p> <p>(1) 曼陀羅 (<i>Datura stramonium</i>)</p> <p>(2) 羅勒 (<i>Ocimum basilicum</i>)</p> <p>(3) 茄屬 (<i>Solanum</i> spp.)</p>	<p>亞洲及太平洋地區</p> <p>(1) 敘利亞</p> <p>(2) 土耳其</p> <p>非洲</p> <p>(3) 卡那利群島 (西班牙屬)</p> <p>(4) 南非</p> <p>歐洲</p> <p>(5) 奧地利</p> <p>(6) 比利時</p> <p>(7) 保加利亞</p> <p>(8) 賽普勒斯</p> <p>(9) 丹麥</p> <p>(10) 法國</p> <p>(11) 德國</p> <p>(12) 希臘</p> <p>(13) 匈牙利</p> <p>(14) 愛爾蘭</p> <p>(15) 義大利</p> <p>(16) 立陶宛</p> <p>(17) 荷蘭</p> <p>(18) 波蘭</p> <p>(19) 西班牙</p> <p>(20) 瑞士</p> <p>(21) 烏克蘭</p> <p>北美</p> <p>(22) 加拿大(英屬</p>	<p>香瓜茄嵌紋病毒(<i>Pepino mosaic virus</i> (<i>Potexvirus</i>))</p>	<p>1、繁殖用生植株應檢附輸出國植物檢疫機關(構)簽發之植物檢疫證明書,證明經田間檢疫或於輸出前經實驗室檢測未染香瓜茄嵌紋病毒,否則銷燬或退運。</p> <p>2、繁殖用種子應檢附輸出國植物檢疫機關(構)簽發之植物檢疫證明書,證明下列條件之一,否則銷燬或退運：</p> <p>(1)經田間檢疫未染香瓜茄嵌紋病毒,並於採種後經磷酸三鈉檢疫處理。</p> <p>(2)輸出前經實驗室檢測未染香瓜茄嵌紋病毒。</p>	<p>一、配合增訂之乙、十七點「繁殖用生植株」之定義,「種子」已涵括於「繁殖用生植株」內,爰酌修文字,修正前後本項規範之植物部位範圍未變動。</p> <p>二、配合乙、第十六點及其附件「茄屬及番椒屬種子輸入檢疫條件」,並依查詢科學文獻結果,修正國家或地區欄,增列七個發生國家,另序號配合調整。</p>

	<p>(23) 波蘭 (24) 西班牙 (25) 瑞士 (26) 烏克蘭 (27) 塞爾維亞 (28) 英國</p> <p>北美</p> <p>(29) 加拿大(英屬哥倫比亞省及安大略省) (30) 墨西哥 (31) 美國(阿拉巴馬州、亞利桑那州、加利福尼亞州、科羅拉多州、佛羅里達州、馬里蘭州、明尼蘇達州、奧克拉荷馬州、德克薩斯州)</p> <p>中南美</p> <p>(32) 智利 (33) 厄瓜多 (34) 秘魯</p>				<p>哥倫比亞省及安大略省) (23) 墨西哥 (24) 美國(阿拉巴馬州、亞利桑那州、加利福尼亞州、科羅拉多州、佛羅里達州、馬里蘭州、明尼蘇達州、奧克拉荷馬州、德克薩斯州) 中南美 (25) 智利 (26) 厄瓜多 (27) 秘魯</p>			
<p>43.下列繁殖用種子： (1) 番椒屬 (<i>Capsicum</i> spp.) (2) 茄屬 (<i>Solanum</i> spp.)</p>	<p>亞洲及太平洋地區</p> <p>(1) 澳大利亞 (2) 巴林 (3) 孟加拉 (4) 中國大陸 (5) 法屬玻里尼西亞 (6) 喬治亞 (7) 印度 (8) 伊朗 (9) 伊拉克 (10) 以色列 (11) 日本 (12) 約旦</p>	<p>番茄黃化捲葉病毒 (<i>Tomato yellow leaf curl virus</i> (Begomovirus))</p>	<p>應檢附輸出國植物檢疫機關(構)簽發之植物檢疫證明書，證明輸出前經實驗室檢測未罹染番茄黃化捲葉病毒，否則銷燬或退運。</p>					<p>一、本項新增。 二、依據防檢局植物防疫檢疫諮議委員會檢疫小組會議決議，新增番茄黃化捲葉病毒(<i>Tomato yellow leaf curl virus</i> (Begomovirus))之檢疫條件，爰增列第四十三項。</p>

	(13) 科威特 (14) 寮國 (15) 黎巴嫩 (16) 尼泊爾 (17) 新喀里多尼亞 (18) 阿曼 (19) 巴基斯坦 (20) 沙烏地阿拉伯 (21) 南韓 (22) 泰國 (23) 土耳其 (24) 阿拉伯聯合大 公國 (25) 葉門 非洲 (26) 阿爾及利亞 (27) 貝南 (28) 布吉納法索 (29) 維德角 (30) 象牙海岸 (31) 埃及 (32) 迦納 (33) 利比亞 (34) 馬利 (35) 模里西斯 (36) 摩洛哥 (37) 奈及利亞 (38) 留尼旺 (39) 塞內加爾 (40) 蘇丹 (41) 坦尚尼亞 (42) 突尼西亞 (43) 烏干達 (44) 亞塞拜然 歐洲 (45) 賽普勒斯 (46) 法國 (47) 希臘 (48) 義大利 (49) 馬爾他				
--	--	--	--	--	--

	(50) 葡萄牙 (51) 西班牙 (52) 瑞士 北美 (53) 墨西哥 (54) 美國 中南美 (55) 巴哈馬 (56) 哥斯大黎加 (57) 古巴 (58) 多米尼克 (59) 多明尼加 (60) 格瑞那達 (61) 瓜地洛普 (62) 瓜地馬拉 (63) 牙買加 (64) 馬丁尼克 (65) 波多黎各 (66) 聖克里斯多福 及尼維斯 (67) 委內瑞拉				
44.下列繁殖用種子： (1) 番椒屬 (<i>Capsicum</i> spp.) (2) 茄屬 (<i>Solanum</i> spp.)	亞洲及太平洋地區 (1) 中國大陸 (2) 伊朗 (3) 以色列 (4) 越南 歐洲 (5) 捷克 (6) 西班牙 北美 (7) 墨西哥 (8) 美國 中南美 (9) 巴西	番茄斑駁嵌紋病毒(<i>Tomato mottle mosaic virus</i>)	應檢附輸出國植物檢疫機關(構)簽發之植物檢疫證明書,證明輸出前經實驗室檢測未罹染番茄斑駁嵌紋病毒,否則銷燬或退運。		一、 <u>本項新增</u> 。 二、依據防檢局植物防疫檢疫諮議委員會檢疫小組會議決議,新增番茄斑駁嵌紋病毒(<i>Tomato mottle mosaic virus</i>)之檢疫條件,爰增列第四十四項。
45.下列繁殖用種子： (1) 番椒 (<i>Capsicum annuum</i>) (2) 茄屬 (<i>Solanum</i> spp.)	亞洲及太平洋地區 (1) 泰國 (2) 越南 歐洲 (3) 荷蘭 北美	番椒小果類病毒(<i>Pepper chat fruit viroid, PCFVd</i>)	應檢附輸出國植物檢疫機關(構)簽發之植物檢疫證明書,證明輸出前經實驗室檢測		一、 <u>本項新增</u> 。 二、依據防檢局植物防疫檢疫諮議委員會檢疫小組會議決議,新增番椒小果類病毒(<i>Pepper chat fruit viroid, PCFVd</i>)之檢疫條件,爰增列第四十五項。

	(4) 加拿大		未罹染番椒小果類病毒，否則銷燬或退運。		
46.下列繁殖用種子： (1) 番椒 (<i>Capsicum annuum</i>) (2) 茄屬 (<i>Solanum</i> spp.)	亞洲及太平洋地區 (1) 泰國 非洲 (2) 迦納 (3) 馬利 歐洲 (4) 比利時 (5) 丹麥 (6) 法國 (7) 德國 (8) 義大利 (9) 荷蘭 (10) 葡萄牙 (11) 英國 北美 (12) 加拿大 (13) 美國 中南美 (14) 哥斯大黎加	鯨魚藤潛隱類病毒 (<i>Columnea latent viroid</i> , CLVd)	應檢附輸出國植物檢疫機關(構)簽發之植物檢疫證明書，證明輸出前經實驗室檢測未罹染鯨魚藤潛隱類病毒，否則銷燬或退運。		一、本項新增。 二、依據防檢局植物防疫檢疫諮議委員會檢疫小組會議決議，新增鯨魚藤潛隱類病毒(<i>Columnea latent viroid</i> , CLVd)之檢疫條件，爰增列第四十六項。
47.下列繁殖用種子： (1) 番椒 (<i>Capsicum annuum</i>) (2) 茄屬 (<i>Solanum</i> spp.)	亞洲及太平洋地區 (1) 印尼 (2) 以色列 (3) 日本 (4) 約旦 (5) 巴勒斯坦 (6) 泰國 (7) 土耳其 非洲 (8) 迦納 (9) 塞內加爾 (10) 突尼西亞 (11) 象牙海岸	番茄莖頂矮化類病毒 (<i>Tomato apical stunt viroid</i> , TASVd)	應檢附輸出國植物檢疫機關(構)簽發之植物檢疫證明書，證明輸出前經實驗室檢測未罹染番茄莖頂矮化類病毒，否則銷燬或退運。		一、本項新增。 二、依據防檢局植物防疫檢疫諮議委員會檢疫小組會議決議，新增番茄莖頂矮化類病毒(<i>Tomato apical stunt viroid</i> , TASVd)之檢疫條件，爰增列第四十七項。

	<p>歐洲</p> <p>(12) 奧地利</p> <p>(13) 比利時</p> <p>(14) 克羅埃西亞</p> <p>(15) 愛沙尼亞</p> <p>(16) 芬蘭</p> <p>(17) 法國</p> <p>(18) 德國</p> <p>(19) 義大利</p> <p>(20) 荷蘭</p> <p>(21) 波蘭</p> <p>(22) 斯洛伐尼亞</p>				
<p>48.下列繁殖用種子：</p> <p>(1) 番椒 (<i>Capsicum annuum</i>)</p> <p>(2) 茄屬 (<i>Solanum</i> spp.)</p>	<p>北美</p> <p>(1) 加拿大</p> <p>(2) 墨西哥</p>	<p>番茄植株雄化類病毒 (<i>Tomato planta macho viroid</i>, TPMVd)</p>	<p>應檢附輸出國植物檢疫機關(構)簽發之植物檢疫證明書，證明輸出前經實驗室檢測未罹染番茄植株雄化類病毒，否則銷燬或退運。</p>		<p>一、本項新增。</p> <p>二、依據防檢局植物防疫檢疫諮議委員會檢疫小組會議決議，新增番茄植株雄化類病 (<i>Tomato planta macho viroid</i>, TPMVd)之檢疫條件，爰增列第四十八項。</p>
<p>49.下列繁殖用種子：</p> <p>(1) 番椒 (<i>Capsicum annuum</i>)</p> <p>(2) 茄屬 (<i>Solanum</i> spp.)</p>	<p>亞洲及太平洋地區</p> <p>(1) 印度</p> <p>歐洲</p> <p>(2) 捷克</p> <p>(3) 法國</p> <p>(4) 芬蘭</p> <p>(5) 斯洛伐尼亞</p> <p>(6) 英國</p> <p>北美</p> <p>(7) 墨西哥</p> <p>(8) 美國</p>	<p>番茄褪綠矮化類病毒 (<i>Tomato chlorotic dwarf viroid</i>, TCDVd)</p>	<p>應檢附輸出國植物檢疫機關(構)簽發之植物檢疫證明書，證明輸出前經實驗室檢測未罹染番茄褪綠矮化類病毒，否則銷燬或退運。</p>		<p>一、本項新增。</p> <p>二、依據防檢局植物防疫檢疫諮議委員會檢疫小組會議決議，新增番茄褪綠矮化類病毒(<i>Tomato chlorotic dwarf viroid</i>, TCDVd)之檢疫條件，爰增列第四十九項。</p>

十、輸入之植物或植物產品經檢疫結果證明有下列有害生物存在，應經適當之檢疫處理，確定該有害生物完全滅除後，始得輸入。無適當之檢疫處理方式可滅除該有害生物時，應予退運或銷燬。行政院農業委員會動植物防疫檢疫局得對下列有害生物之上一分類群 (Taxa) 有害生物採取檢疫處理措施。

病毒類 (467 種)

Abutilon mosaic virus (AbMV) (Geminiviridae, Begomovirus)
Ageratum yellow vein virus (AYVV) (Geminiviridae, Begomovirus)
African cassava mosaic virus (ACMV) (Geminiviridae, Begomovirus)
Agropyron mosaic virus (AgMV) (Potyviridae, Bymovirus)
Alfalfa cryptic virus 1 (ACV-1) (Partitiviridae, alphacryptovirus)
Alfalfa enation virus
Alfalfa mosaic virus (AIMV) (Bromoviridae, alfamovirus)
Alsike clover vein mosaic virus
Alstroemeria mosaic virus (AIMV) (Potyviridae, potyvirus)
Amaranthus leaf mottle virus (AmLMV) (Potyviridae, potyvirus)
American plum line pattern virus (APLPV) (Bromoviridae, Ilarvirus)
Andean potato latent virus (APLV) (Tymovirus)
Andean potato mottle virus (APMoV) (Comoviridae, Comovirus)
Anthoxanthum latent blanching virus (ALBV) (Hordeivirus)
Anthriscus yellows virus (AYV) (Sequiviridae, waikavirus)
Apple chat fruit disease
Apple chlorotic leaf spot virus (ACLSV) (Trichovirus, Flexiviridae)
Apple dimple fruit viroid (ADFVd) (Apscaviroid, Pospiviroidae)
Apple mosaic virus (ApMV) (Bromoviridae, Ilarvirus)
Apple scar skin viroid (ASSVd) (Pospiviroidae, Apscaviroid)
Apple stem grooving virus (ASGV) (Capillovirus, Flexiviridae)
Apple stem pitting virus (ASPV) (Foveavirus, Flexiviridae)
Araujia mosaic virus (ArjMV) (Potyviridae, potyvirus)
Arracacha Virus A (AVA) (Comoviridae, Nepovirus)
Arracacha Virus B (AVB) (Comoviridae, Nepovirus)
Arracacha Virus Y (AVY) (Potyviridae, potyvirus)
Artichoke Italian latent virus (AILV) (Comoviridae, nepovirus)
Artichoke latent virus (ArLV) (Potyviridae, potyvirus)
Artichoke mottled crinkle virus (AMCV) (Tombusviridae, Tombusvirus)
Artichoke yellow ringspot virus (AYRSV) (Comoviridae, Nepovirus)
Asparagus virus 1 (AV-1) (Potyviridae, potyvirus)
Asparagus virus 2 (AV-2) (Bromoviridae, ilarvirus)
Asparagus virus 3 (AV-3) (Potexvirus)
Banana bunchy top virus (BBTV) (Nanovirus)
Barley mild mosaic virus (BaMMV) (Potyviridae, bymovirus)
Barley mosaic virus (BBV)
Barley yellow dwarf virus-GPV (BYDV-GPV) (Luteovirus)
Barley yellow mosaic virus (BaYMV) (Potyviridae, Bymovirus)
Barley yellow striate mosaic virus (BYSMV) (Rhabdoviridae, Cytorhabdovirus)
Bean calico mosaic virus (BCaMV) (Geminiviridae, Begomovirus)
Bean common mosaic virus (Potyviridae, potyvirus)
Bean golden mosaic virus (BGMV) (Geminiviridae, Begomovirus)
Bean leaf roll virus (BLRV) (Luteoviridae, luteovirus)
Bean mild mosaic virus (BMMV) (Tombusviridae, carmovirus)

十、輸入之植物或植物產品經檢疫結果證明有下列有害生物存在，應經適當之檢疫處理，確定該有害生物完全滅除後，始得輸入。無適當之檢疫處理方式可滅除該有害生物時，應予退運或銷燬。行政院農業委員會動植物防疫檢疫局得對下列有害生物之上一分類群 (Taxa) 有害生物採取檢疫處理措施。

病毒類 (474 種)

Abutilon mosaic virus (AbMV) (Geminiviridae, Begomovirus)
Ageratum yellow vein virus (AYVV) (Geminiviridae, Begomovirus)
African cassava mosaic virus (ACMV) (Geminiviridae, Begomovirus)
Agropyron mosaic virus (AgMV) (Potyviridae, Bymovirus)
Alfalfa cryptic virus 1 (ACV-1) (Partitiviridae, alphacryptovirus)
Alfalfa enation virus
Alfalfa mosaic virus (AIMV) (Bromoviridae, alfamovirus)
Alsike clover vein mosaic virus
Alstroemeria mosaic virus (AIMV) (Potyviridae, potyvirus)
Amaranthus leaf mottle virus (AmLMV) (Potyviridae, potyvirus)
American plum line pattern virus (APLPV) (Bromoviridae, Ilarvirus)
Andean potato latent virus (APLV) (Tymovirus)
Andean potato mottle virus (APMoV) (Comoviridae, Comovirus)
Anthoxanthum latent blanching virus (ALBV) (Hordeivirus)
Anthriscus yellows virus (AYV) (Sequiviridae, waikavirus)
Apple chat fruit disease
Apple chlorotic leaf spot virus (ACLSV) (Trichovirus, Flexiviridae)
Apple dimple fruit viroid (ADFVd) (Apscaviroid, Pospiviroidae)
Apple mosaic virus (ApMV) (Bromoviridae, Ilarvirus)
Apple scar skin viroid (ASSVd) (Pospiviroidae, Apscaviroid)
Apple stem grooving virus (ASGV) (Capillovirus, Flexiviridae)
Apple stem pitting virus (ASPV) (Foveavirus, Flexiviridae)
Araujia mosaic virus (ArjMV) (Potyviridae, potyvirus)
Arracacha Virus A (AVA) (Comoviridae, Nepovirus)
Arracacha Virus B (AVB) (Comoviridae, Nepovirus)
Arracacha Virus Y (AVY) (Potyviridae, potyvirus)
Artichoke Italian latent virus (AILV) (Comoviridae, nepovirus)
Artichoke latent virus (ArLV) (Potyviridae, potyvirus)
Artichoke mottled crinkle virus (AMCV) (Tombusviridae, Tombusvirus)
Artichoke yellow ringspot virus (AYRSV) (Comoviridae, Nepovirus)
Asparagus virus 1 (AV-1) (Potyviridae, potyvirus)
Asparagus virus 2 (AV-2) (Bromoviridae, ilarvirus)
Asparagus virus 3 (AV-3) (Potexvirus)
Banana bunchy top virus (BBTV) (Nanovirus)
Barley mild mosaic virus (BaMMV) (Potyviridae, bymovirus)
Barley mosaic virus (BBV)
Barley yellow dwarf virus-GPV (BYDV-GPV) (Luteovirus)
Barley yellow mosaic virus (BaYMV) (Potyviridae, Bymovirus)
Barley yellow striate mosaic virus (BYSMV) (Rhabdoviridae, Cytorhabdovirus)
Bean calico mosaic virus (BCaMV) (Geminiviridae, Begomovirus)
Bean common mosaic virus (Potyviridae, potyvirus)
Bean golden mosaic virus (BGMV) (Geminiviridae, Begomovirus)
Bean leaf roll virus (BLRV) (Luteoviridae, luteovirus)
Bean mild mosaic virus (BMMV) (Tombusviridae, carmovirus)

配合魚藤潛隱類病毒 (*Columnnea latent viroid*, CLVd)、番椒小果類病毒 (*Pepper chat fruit viroid*, PCFVd)、番茄莖頂矮化類病毒 (*Tomato pical stunt viroid*, TASVd)、番茄褪綠矮化類病毒 (*Tomato chlorotic dwarf viroid*, TCDVd)、番茄斑駁嵌紋病毒 (*Tomato mottle mosaic virus*, ToMMV)、番茄植株雄化類病毒 (*Tomato planta macho viroid*, TPMVd) 及番茄黃化捲葉病毒 (*Tomato yellow leaf curl virus*, TYLCV) 於第一點訂定檢疫條件，爰自清單刪除。

<p><i>Bean pod mottle virus (BPMV) (Comoviridae, comovirus)</i> <i>Bean rugose mosaic virus (BRMV) (Comoviridae, comovirus)</i> <i>Bean southern mosaic virus (SBMV) (sobemovirus)</i> <i>Bean yellow vein banding virus (BYVBV) (Umbravirus)</i> <i>Beet distortion mosaic virus</i> <i>Beet mild yellowing virus (BMYV) (Luteoviridae, luteovirus)</i> <i>Beet mosaic virus (BtMV) (Potyviridae, potyvirus)</i> <i>Beet soil-borne virus (BSBV) (Pomovirus)</i> <i>Beet western yellows virus (BWYV) (Luteoviridae, luteovirus)</i> <i>Beet yellows virus (BYV) (Closteroviridae, closterovirus)</i> <i>Beet yellow stunt virus (BYSV) (Closteroviridae, closterovirus)</i> <i>Beet curly top virus (BCTV) (Geminiviridae, Curtomovirus)</i> <i>Beet leaf curl virus (BLCV) (Rhabdoviridae)</i> <i>Beet necrotic yellow vein virus (BNYVV) (Benyvirus)</i> <i>Belladonna mottle virus (BeMV) (Tymovirus)</i> <i>Bermuda grass etched-line virus (BELV) (Marafivirus)</i> <i>Bhendi yellow vein mosaic virus (BYVMV) (Geminiviridae, Begomovirus)</i> <i>Bidens mosaic virus (BiMV) (Potyviridae, potyvirus)</i> <i>Bidens mottle virus (BiMoV) (Potyviridae, potyvirus)</i> <i>Black raspberry latent ilarvirus</i> <i>Black raspberry necrosis virus (BRNV)</i> <i>Blackcurrant reversion associated virus (BRAV) (Comoviridae, Nepovirus)</i> <i>Blueberry leaf mottle virus (BLMoV) (Comoviridae, Nepovirus)</i> <i>Blueberry necrotic shock virus (BIShV) (ilarvirus)</i> <i>Blueberry red ringspot virus (BRRV) (Caulimoviridae, caulimovirus)</i> <i>Blueberry scorch virus (BBScV) (carlavirus)</i> <i>Blueberry shoestring virus (BSSV) (sobemovirus)</i> <i>Broad bean B virus</i> <i>Broad bean necrosis virus (BBNV) (Pomovirus)</i> <i>Broad bean severe chlorosis virus (Closteroviridae, closterovirus)</i> <i>Broad bean stain virus (BBSV) (Comoviridae, comovirus)</i> <i>Broad bean true mosaic virus (BBTMV) (Comoviridae, comovirus)</i> <i>Broad bean wilt virusd (BBWV) (Comoviridae, fabavirus)</i> <i>Broad bean yellow ringspot virus</i> <i>Broad bean yellow vein virus (Rhabdoviridae, cytorhabdovirus)</i> <i>Broccoli necrotic yellows virus (BNYV) (Rhabdoviridae, cytorhabdovirus)</i> <i>Brome mosaic virus (BMV) (Bromoviridae, bromovirus)</i> <i>Brome streak mosaic virus (BrStMV) (Potyviridae, Tritimovirus)</i> <i>Burdock mosaic virus</i> <i>Burdock mottle virus</i> <i>Burdock yellows virus (BuYV) (Closteroviridae, closterovirus)</i> <i>Cacao necrosis virus (CNV) (Comoviridae, nepovirus)</i> <i>Cacao swollen shoot virus (CSSV) (Caulimoviridae, badnavirus)</i> <i>Cacao yellow mosaic virus (CYMV) (Tymovirus)</i> <i>Cactus virus (CV-2) (Carlavirus)</i> <i>Canna yellow mottle VIRUS (CaYMV) (Caulimoviridae, badnavirus)</i> <i>Caper latent virus (CapLV) (Carlavirus)</i> <i>Cardamine chlorotic fleck virus (CCFV) (Tombusviridae, carmovirus)</i> <i>Cardamom mosaic virus (CdMV) (Potyviridae, potyvirus)</i> <i>Carnation cryptic virus 1 (CCV-1) (Partitiviridae, alphacryptovirus)</i> <i>Carnation etched ring virus (CERV) (Cualimoviridae, caulimovirus)</i> <i>Carnation Italian ringspot virus (CIRV) (Tombusviridae, tombusvirus)</i></p>	<p><i>Bean pod mottle virus (BPMV) (Comoviridae, comovirus)</i> <i>Bean rugose mosaic virus (BRMV) (Comoviridae, comovirus)</i> <i>Bean southern mosaic virus (SBMV) (sobemovirus)</i> <i>Bean yellow vein banding virus (BYVBV) (Umbravirus)</i> <i>Beet distortion mosaic virus</i> <i>Beet mild yellowing virus (BMYV) (Luteoviridae, luteovirus)</i> <i>Beet mosaic virus (BtMV) (Potyviridae, potyvirus)</i> <i>Beet soil-borne virus (BSBV) (Pomovirus)</i> <i>Beet western yellows virus (BWYV) (Luteoviridae, luteovirus)</i> <i>Beet yellows virus (BYV) (Closteroviridae, closterovirus)</i> <i>Beet yellow stunt virus (BYSV) (Closteroviridae, closterovirus)</i> <i>Beet curly top virus (BCTV) (Geminiviridae, Curtomovirus)</i> <i>Beet leaf curl virus (BLCV) (Rhabdoviridae)</i> <i>Beet necrotic yellow vein virus (BNYVV) (Benyvirus)</i> <i>Belladonna mottle virus (BeMV) (Tymovirus)</i> <i>Bermuda grass etched-line virus (BELV) (Marafivirus)</i> <i>Bhendi yellow vein mosaic virus (BYVMV) (Geminiviridae, Begomovirus)</i> <i>Bidens mosaic virus (BiMV) (Potyviridae, potyvirus)</i> <i>Bidens mottle virus (BiMoV) (Potyviridae, potyvirus)</i> <i>Black raspberry latent ilarvirus</i> <i>Black raspberry necrosis virus (BRNV)</i> <i>Blackcurrant reversion associated virus (BRAV) (Comoviridae, Nepovirus)</i> <i>Blueberry leaf mottle virus (BLMoV) (Comoviridae, Nepovirus)</i> <i>Blueberry necrotic shock virus (BIShV) (ilarvirus)</i> <i>Blueberry red ringspot virus (BRRV) (Caulimoviridae, caulimovirus)</i> <i>Blueberry scorch virus (BBScV) (carlavirus)</i> <i>Blueberry shoestring virus (BSSV) (sobemovirus)</i> <i>Broad bean B virus</i> <i>Broad bean necrosis virus (BBNV) (Pomovirus)</i> <i>Broad bean severe chlorosis virus (Closteroviridae, closterovirus)</i> <i>Broad bean stain virus (BBSV) (Comoviridae, comovirus)</i> <i>Broad bean true mosaic virus (BBTMV) (Comoviridae, comovirus)</i> <i>Broad bean wilt virusd (BBWV) (Comoviridae, fabavirus)</i> <i>Broad bean yellow ringspot virus</i> <i>Broad bean yellow vein virus (Rhabdoviridae, cytorhabdovirus)</i> <i>Broccoli necrotic yellows virus (BNYV) (Rhabdoviridae, cytorhabdovirus)</i> <i>Brome mosaic virus (BMV) (Bromoviridae, bromovirus)</i> <i>Brome streak mosaic virus (BrStMV) (Potyviridae, Tritimovirus)</i> <i>Burdock mosaic virus</i> <i>Burdock mottle virus</i> <i>Burdock yellows virus (BuYV) (Closteroviridae, closterovirus)</i> <i>Cacao necrosis virus (CNV) (Comoviridae, nepovirus)</i> <i>Cacao swollen shoot virus (CSSV) (Caulimoviridae, badnavirus)</i> <i>Cacao yellow mosaic virus (CYMV) (Tymovirus)</i> <i>Cactus virus (CV-2) (Carlavirus)</i> <i>Canna yellow mottle VIRUS (CaYMV) (Caulimoviridae, badnavirus)</i> <i>Caper latent virus (CapLV) (Carlavirus)</i> <i>Cardamine chlorotic fleck virus (CCFV) (Tombusviridae, carmovirus)</i> <i>Cardamom mosaic virus (CdMV) (Potyviridae, potyvirus)</i> <i>Carnation cryptic virus 1 (CCV-1) (Partitiviridae, alphacryptovirus)</i> <i>Carnation etched ring virus (CERV) (Cualimoviridae, caulimovirus)</i> <i>Carnation Italian ringspot virus (CIRV) (Tombusviridae, tombusvirus)</i></p>	
--	--	--

<p><i>Carnation latent virus (CLV) (Carlavirus)</i> <i>Carnation mottle virus (CarMV) (Carmovirus)</i> <i>Carnation necrotic fleck virus (CNFV) (Clostreoviridae, closterovirus)</i> <i>Carnation ringspot virus (CRSV) (Tombusviridae, dianthovirus)</i> <i>Carnation vein mottle virus (CVMoV) (Potyviridae, potyvirus)</i> <i>Carrot latent virus (CtLV) (Rhabdoviridae, nucleorhabdovirus)</i> <i>Carrot mottle mimic virus (CmoMV) (Umbravirus)</i> <i>Carrot mottle virus (CMoV) (Umbravirus)</i> <i>Carrot red leaf virus (CtRLV) (Luteoviridae, luteovirus)</i> <i>Carrot temperate virus 1 (CteV-1) (Partitiviridae, alphacryptovirus)</i> <i>Carrot temperate virus 2 (CteV-2) (Partitiviridae, betacryptovirus)</i> <i>Carrot temperate virus 3 (CteV-3) (Partitiviridae, alphacryptovirus)</i> <i>Carrot temperate virus 4 (CteV-4) (Partitiviridae, alphacryptovirus)</i> <i>Carrot thin leaf virus (CTLV) (Potyviridae, potyvirus)</i> <i>Cowpea chlorotic mottle virus (CCMV) (Bromoviridae, Bromovirus)</i> <i>Cowpea golden mosaic virus (CPGMV) (Geminiviridae, Begomovirus)</i> <i>Cowpea green vein-banding virus (CGVBV) (Potyviridae, Potyvirus)</i> <i>Cowpea mosaic virus (CPMV) (Comoviridae, Comovirus)</i> <i>Cowpea rugose mosaic virus (CPRMV) (Potyviridae, Potyvirus)</i> <i>Cowpea severe mosaic virus (CPSMV) (Comoviridae, Comovirus)</i> <i>Cowpea stunt virus (Luteovirus)</i> <i>Cassava African mosaic virus (ACMV) (Geminiviridae, bigeminivirus)</i> <i>Cassava green mottle virus (CaGMV) (Comoviridae, nepovirus)</i> <i>Cassava Indian mosaic virus (ICMV) (Geminiviridae, bigeminivirus)</i> <i>Cassava Ivorian bacilliform virus (CsIBV) (Ourmiavirus)</i> <i>Cassava virus X (CsVX) (Potexvirus)</i> <i>Cassia mosaic virus</i> <i>Cassia ringspot virus</i> <i>Cassia yellow blotch virus (CYBV) (Bromoviridae, bromovirus)</i> <i>Cassia yellow spot virus (CasYSV) (Potyviridae, potyvirus)</i> <i>Cauliflower mosaic virus (CaMV) (Caulimoviridae, caulimovirus)</i> <i>Celery mosaic virus (CeMV) (Potyviridae, potyvirus)</i> <i>Cereal northern mosaic virus (NCMV) (cytorhabdovirus)</i> <i>Cassava brown streak virus (CsBSV) (Potyviridae, Ipomovirus)</i> <i>Cassava common mosaic virus (CsCMV) (Potexvirus)</i> <i>Cereal chlorotic mosaic virus (CCMoV) (Rhabdoviridae)</i> <i>Cherry green ring mottle virus (CGRMV) (Foveavirus)</i> <i>Cherry leaf roll virus (CLRV) (Comoviridae, Nepovirus)</i> <i>Cherry mottle leaf virus (CMLV) (Trichovirus)</i> <i>Cherry rasp leaf virus (CRLV) (Comoviridae, Nepovirus)</i> <i>Cherry virus A (CVA) (Capillovirus)</i> <i>Chickpea bushy dwarf virus (CpBDV) (Potyviridae, potyvirus)</i> <i>Chickpea distortion mosaic virus (Potyviridae, potyvirus)</i> <i>Chicory virus (ChVX) (potexvirus)</i> <i>Chicory yellow mottle virus (ChYMV) (Comoviridae, nepovirus)</i> <i>Chino del tomate virus (CdTV) (Geminiviridae, bigeminivirus)</i> <i>Chloris striate mosaic virus (CSMV) (Geminiviridae, monogeminivirus)</i> <i>Chrysanthemum virus B (CVB) (Carlavirus)</i> <i>Citrus blight disease</i> <i>Citrus leprosis virus (CiLV) (Rhabdoviridae)</i> <i>Citrus yellow mosaic virus (CiYMV) (Caulimoviridae, Badnavirus)</i> <i>Citrus leaf rugose virus (CiLRV) (Rhabdoviridae, ilarvirus)</i></p>	<p><i>Carnation latent virus (CLV) (Carlavirus)</i> <i>Carnation mottle virus (CarMV) (Carmovirus)</i> <i>Carnation necrotic fleck virus (CNFV) (Clostreoviridae, closterovirus)</i> <i>Carnation ringspot virus (CRSV) (Tombusviridae, dianthovirus)</i> <i>Carnation vein mottle virus (CVMoV) (Potyviridae, potyvirus)</i> <i>Carrot latent virus (CtLV) (Rhabdoviridae, nucleorhabdovirus)</i> <i>Carrot mottle mimic virus (CmoMV) (Umbravirus)</i> <i>Carrot mottle virus (CMoV) (Umbravirus)</i> <i>Carrot red leaf virus (CtRLV) (Luteoviridae, luteovirus)</i> <i>Carrot temperate virus 1 (CteV-1) (Partitiviridae, alphacryptovirus)</i> <i>Carrot temperate virus 2 (CteV-2) (Partitiviridae, betacryptovirus)</i> <i>Carrot temperate virus 3 (CteV-3) (Partitiviridae, alphacryptovirus)</i> <i>Carrot temperate virus 4 (CteV-4) (Partitiviridae, alphacryptovirus)</i> <i>Carrot thin leaf virus (CTLV) (Potyviridae, potyvirus)</i> <i>Cowpea chlorotic mottle virus (CCMV) (Bromoviridae, Bromovirus)</i> <i>Cowpea golden mosaic virus (CPGMV) (Geminiviridae, Begomovirus)</i> <i>Cowpea green vein-banding virus (CGVBV) (Potyviridae, Potyvirus)</i> <i>Cowpea mosaic virus (CPMV) (Comoviridae, Comovirus)</i> <i>Cowpea rugose mosaic virus (CPRMV) (Potyviridae, Potyvirus)</i> <i>Cowpea severe mosaic virus (CPSMV) (Comoviridae, Comovirus)</i> <i>Cowpea stunt virus (Luteovirus)</i> <i>Cassava African mosaic virus (ACMV) (Geminiviridae, bigeminivirus)</i> <i>Cassava green mottle virus (CaGMV) (Comoviridae, nepovirus)</i> <i>Cassava Indian mosaic virus (ICMV) (Geminiviridae, bigeminivirus)</i> <i>Cassava Ivorian bacilliform virus (CsIBV) (Ourmiavirus)</i> <i>Cassava virus X (CsVX) (Potexvirus)</i> <i>Cassia mosaic virus</i> <i>Cassia ringspot virus</i> <i>Cassia yellow blotch virus (CYBV) (Bromoviridae, bromovirus)</i> <i>Cassia yellow spot virus (CasYSV) (Potyviridae, potyvirus)</i> <i>Cauliflower mosaic virus (CaMV) (Caulimoviridae, caulimovirus)</i> <i>Celery mosaic virus (CeMV) (Potyviridae, potyvirus)</i> <i>Cereal northern mosaic virus (NCMV) (cytorhabdovirus)</i> <i>Cassava brown streak virus (CsBSV) (Potyviridae, Ipomovirus)</i> <i>Cassava common mosaic virus (CsCMV) (Potexvirus)</i> <i>Cereal chlorotic mosaic virus (CCMoV) (Rhabdoviridae)</i> <i>Cherry green ring mottle virus (CGRMV) (Foveavirus)</i> <i>Cherry leaf roll virus (CLRV) (Comoviridae, Nepovirus)</i> <i>Cherry mottle leaf virus (CMLV) (Trichovirus)</i> <i>Cherry rasp leaf virus (CRLV) (Comoviridae, Nepovirus)</i> <i>Cherry virus A (CVA) (Capillovirus)</i> <i>Chickpea bushy dwarf virus (CpBDV) (Potyviridae, potyvirus)</i> <i>Chickpea distortion mosaic virus (Potyviridae, potyvirus)</i> <i>Chicory virus (ChVX) (potexvirus)</i> <i>Chicory yellow mottle virus (ChYMV) (Comoviridae, nepovirus)</i> <i>Chino del tomate virus (CdTV) (Geminiviridae, bigeminivirus)</i> <i>Chloris striate mosaic virus (CSMV) (Geminiviridae, monogeminivirus)</i> <i>Chrysanthemum virus B (CVB) (Carlavirus)</i> <i>Citrus blight disease</i> <i>Citrus leprosis virus (CiLV) (Rhabdoviridae)</i> <i>Citrus yellow mosaic virus (CiYMV) (Caulimoviridae, Badnavirus)</i> <i>Citrus leaf rugose virus (CiLRV) (Rhabdoviridae, ilarvirus)</i></p>	
---	---	--

<p><i>Citrus ringspot virus</i> <i>Citrus sudden death-associated virus</i> (CSD- associated virus) (<i>Marafivirus</i>, <i>Tymoviridae</i>) <i>Citrus variegation virus</i> (CVV) (<i>Bromoviridae</i>, <i>ilarvirus</i>) <i>Clitoria yellow vein virus</i> (CYVV) (<i>Tymovirus</i>) <i>Clover mild mosaic virus</i> <i>Clover wound tumor virus</i> (WTV) (<i>Phytoreovirus</i>) <i>Clover yellow mosaic virus</i> (CIYMV) (<i>potexvirus</i>) <i>Clover yellows virus</i> (CYV) (<i>Closteroviridae</i>, <i>closterovirus</i>) <i>Clover yellow vein virus</i> (CIYVV) (<i>Potyviridae</i>, <i>potyvirus</i>) <i>Cocksfoot mottle virus</i> (CoMV) (<i>Sobemovirus</i>) <i>Coconut foliar decay virus</i> (CFDV) (<i>Nanavirus</i>) <i>Commelina diffusa virus</i> (<i>Potyviridae</i>, <i>potyvirus</i>) <i>Commelina mosaic virus</i> (ComMV) (<i>Potyviridae</i>, <i>potyvirus</i>) <i>Commelina virus X</i> (ComVX) (<i>potexvirus</i>) <i>Commelina yellow mottle virus</i> (ComYMV) (<i>Caulimoviridae</i>, <i>badnavirus</i>)</p> <p><i>Cocksfoot mild mosaic virus</i> (CoMV) (<i>Sobemovirus</i>) <i>Cocoa mottle leaf virus</i> <i>Cocoa necrosis virus</i> (CoNV) (<i>Comoviridae</i>, <i>Nepovirus</i>) <i>Cocoa swollen shoot virus</i> (CSSV) (<i>Caulimoviridae</i>, <i>Badnavirus</i>) <i>Cocoa yellow mosaic virus</i> <i>Colombian datura virus</i> (CDV) (<i>Potyviridae</i>, <i>Potyvirus</i>) <i>Coriander feathery red vein virus</i> (CFRVV) (<i>Rhabdoviridae</i>, <i>nucleorhabdovirus</i>) <i>Cotton leaf curl virus</i> (CLCuV) (<i>Geminiviridae</i>, <i>Begomovirus</i>) <i>Cotton leaf crumple virus</i> (CLCrV) (<i>Geminiviridae</i>, <i>bigeminivirus</i>) <i>Cowpea mild mottle virus</i> (CPMMV) (<i>Carlavirus</i>) <i>Crimson clover latent virus</i> (CCLV) (<i>Comoviridae</i>, <i>nepovirus</i>) <i>Crinum mosaic virus</i> (CriMV) (<i>Potyviridae</i>, <i>potyvirus</i>) <i>Croton yellow vein mosaic virus</i> (CYVMV) (<i>Geminiviridae</i>, <i>bigeminivirus</i>) <i>Cucumber green mottle mosaic virus</i> (CGMMV) (<i>Tobamovirus</i>) <i>Cucumber leaf spot virus</i> (CLSV) (<i>Tombusviridae</i>, <i>carmovirus</i>) <i>Cucumber mosaic virus subgroup II</i> (CMV-II) (<i>Bromoviridae</i>, <i>Cucumovirus</i>) <i>Cucumber necrosis virus</i> (CuNV) (<i>Tombusviridae</i>, <i>tombusvirus</i>) <i>Cucumber soil-borne virus</i> (CuSBV) (<i>Tombusviridae</i>, <i>carmovirus</i>) <i>Cucumber vein yellowing virus</i> (CVYV) <i>Cucurbit chlorotic yellows virus</i> (CCYC) (<i>Closteroviridae</i>, <i>Crinivirus</i>) <i>Cycas necrotic stunt virus</i> (CNSV) (<i>Comoviridae</i>, <i>nepovirus</i>) <i>Cymbidium ringspot virus</i> (CymRSV) (<i>Tombusviridae</i>, <i>tombusvirus</i>) <i>Cynodon chlorotic streak virus</i> <i>Cynosurus mottle virus</i> (CnMoV) (<i>Sobemovirus</i>) <i>Dahlia mosaic virus</i> (DMV) (<i>Caulimoviridae</i>, <i>caulimovirus</i>) <i>Dandelion latent virus</i> (DaLV) (<i>Carlavirus</i>) <i>Dandelion yellow mosaic virus</i> (DaYMV) (<i>Sequiviridae</i>, <i>sequivirus</i>) <i>Daphne virus X</i> (DVX) (<i>Potexvirus</i>) <i>Daphne virus Y</i> (DVY) (<i>Potyviridae</i>, <i>potyvirus</i>) <i>Datura Colombian virus</i> (CDV) (<i>Potyviridae</i>, <i>potyvirus</i>) <i>Datura distortion mosaic virus</i> (DDMV) (<i>Potyviridae</i>, <i>potyvirus</i>) <i>Datura necrosis virus</i> (DNV) (<i>Potyviridae</i>, <i>potyvirus</i>) <i>Datura shoestring virus</i> (DSTV) (<i>Potyviridae</i>, <i>potyvirus</i>) <i>Datura yellow vein virus</i> (DYVV) (<i>Rhabdoviridae</i>, <i>nucleorhabdovirus</i>) <i>Datura distortion virus</i></p>	<p><i>Citrus ringspot virus</i> <i>Citrus sudden death-associated virus</i> (CSD- associated virus) (<i>Marafivirus</i>, <i>Tymoviridae</i>) <i>Citrus variegation virus</i> (CVV) (<i>Bromoviridae</i>, <i>ilarvirus</i>) <i>Clitoria yellow vein virus</i> (CYVV) (<i>Tymovirus</i>) <i>Clover mild mosaic virus</i> <i>Clover wound tumor virus</i> (WTV) (<i>Phytoreovirus</i>) <i>Clover yellow mosaic virus</i> (CIYMV) (<i>potexvirus</i>) <i>Clover yellows virus</i> (CYV) (<i>Closteroviridae</i>, <i>closterovirus</i>) <i>Clover yellow vein virus</i> (CIYVV) (<i>Potyviridae</i>, <i>potyvirus</i>) <i>Cocksfoot mottle virus</i> (CoMV) (<i>Sobemovirus</i>) <i>Coconut foliar decay virus</i> (CFDV) (<i>Nanavirus</i>) <i>Commelina diffusa virus</i> (<i>Potyviridae</i>, <i>potyvirus</i>) <i>Commelina mosaic virus</i> (ComMV) (<i>Potyviridae</i>, <i>potyvirus</i>) <i>Commelina virus X</i> (ComVX) (<i>potexvirus</i>) <i>Commelina yellow mottle virus</i> (ComYMV) (<i>Caulimoviridae</i>, <i>badnavirus</i>) <u><i>Columnea latent viroid</i> (CLVd) (<i>Pospiviroidae</i>, <i>Pospiviroid</i>)</u> <i>Cocksfoot mild mosaic virus</i> (CoMV) (<i>Sobemovirus</i>) <i>Cocoa mottle leaf virus</i> <i>Cocoa necrosis virus</i> (CoNV) (<i>Comoviridae</i>, <i>Nepovirus</i>) <i>Cocoa swollen shoot virus</i> (CSSV) (<i>Caulimoviridae</i>, <i>Badnavirus</i>) <i>Cocoa yellow mosaic virus</i> <i>Colombian datura virus</i> (CDV) (<i>Potyviridae</i>, <i>Potyvirus</i>) <i>Coriander feathery red vein virus</i> (CFRVV) (<i>Rhabdoviridae</i>, <i>nucleorhabdovirus</i>) <i>Cotton leaf curl virus</i> (CLCuV) (<i>Geminiviridae</i>, <i>Begomovirus</i>) <i>Cotton leaf crumple virus</i> (CLCrV) (<i>Geminiviridae</i>, <i>bigeminivirus</i>) <i>Cowpea mild mottle virus</i> (CPMMV) (<i>Carlavirus</i>) <i>Crimson clover latent virus</i> (CCLV) (<i>Comoviridae</i>, <i>nepovirus</i>) <i>Crinum mosaic virus</i> (CriMV) (<i>Potyviridae</i>, <i>potyvirus</i>) <i>Croton yellow vein mosaic virus</i> (CYVMV) (<i>Geminiviridae</i>, <i>bigeminivirus</i>) <i>Cucumber green mottle mosaic virus</i> (CGMMV) (<i>Tobamovirus</i>) <i>Cucumber leaf spot virus</i> (CLSV) (<i>Tombusviridae</i>, <i>carmovirus</i>) <i>Cucumber mosaic virus subgroup II</i> (CMV-II) (<i>Bromoviridae</i>, <i>Cucumovirus</i>) <i>Cucumber necrosis virus</i> (CuNV) (<i>Tombusviridae</i>, <i>tombusvirus</i>) <i>Cucumber soil-borne virus</i> (CuSBV) (<i>Tombusviridae</i>, <i>carmovirus</i>) <i>Cucumber vein yellowing virus</i> (CVYV) <i>Cucurbit chlorotic yellows virus</i> (CCYC) (<i>Closteroviridae</i>, <i>Crinivirus</i>) <i>Cycas necrotic stunt virus</i> (CNSV) (<i>Comoviridae</i>, <i>nepovirus</i>) <i>Cymbidium ringspot virus</i> (CymRSV) (<i>Tombusviridae</i>, <i>tombusvirus</i>) <i>Cynodon chlorotic streak virus</i> <i>Cynosurus mottle virus</i> (CnMoV) (<i>Sobemovirus</i>) <i>Dahlia mosaic virus</i> (DMV) (<i>Caulimoviridae</i>, <i>caulimovirus</i>) <i>Dandelion latent virus</i> (DaLV) (<i>Carlavirus</i>) <i>Dandelion yellow mosaic virus</i> (DaYMV) (<i>Sequiviridae</i>, <i>sequivirus</i>) <i>Daphne virus X</i> (DVX) (<i>Potexvirus</i>) <i>Daphne virus Y</i> (DVY) (<i>Potyviridae</i>, <i>potyvirus</i>) <i>Datura Colombian virus</i> (CDV) (<i>Potyviridae</i>, <i>potyvirus</i>) <i>Datura distortion mosaic virus</i> (DDMV) (<i>Potyviridae</i>, <i>potyvirus</i>) <i>Datura necrosis virus</i> (DNV) (<i>Potyviridae</i>, <i>potyvirus</i>) <i>Datura shoestring virus</i> (DSTV) (<i>Potyviridae</i>, <i>potyvirus</i>) <i>Datura yellow vein virus</i> (DYVV) (<i>Rhabdoviridae</i>, <i>nucleorhabdovirus</i>) <i>Datura distortion virus</i></p>	
---	---	--

<p><i>Datura enation mosaic virus</i> <i>Dendrobium mosaic virus</i> (DeMV) (Potyviridae, potyvirus) <i>Desmodium mosaic virus</i> (DesMV) (Potyviridae, potyvirus) <i>Desmodium yellow mottle virus</i> (DYMV) (Tymovirus) <i>Digitaria streak virus</i> (DSV) (Geminiviridae, monogeminivirus) <i>Digitaria striate mosaic</i> (DiSMV) (Geminiviridae, monogeminivirus) <i>Dioscorea alata virus</i> (Potyviridae, potyvirus) <i>Dioscorea bacilliform virus</i> (DBV) (Caulimoviridae, badnavirus) <i>Dioscorea green banding mosaic virus</i> (Potyviridae, potyvirus) <i>Dodonaea yellows-associated virus</i> <i>Dolichos yellow mosaic virus</i> (DoMYV) (Geminiviridae, bigeminivirus) <i>Dulcamara mottle virus</i> (DuMV) (Tymovirus) <i>Eggplant green mosaic virus</i> (EGMV) (Potyviridae, potyvirus) <i>Eggplant mottled dwarf virus</i> (EMDV) (Rhabdoviridae, nucleorhabdovirus) <i>Elderberry virus</i> (EV) (Carlavirus 1) <i>Epirus cherry virus</i> (EpCV) (Ourmiavirus) <i>Erysimum latent virus</i> (ErLV) (Tymovirus) <i>Echinochloa ragged stunt virus</i> (ERSV) (Reoviridae, Oryzavirus) <i>Elm mottle virus</i> (EMoV) (Bromoviridae, Ilarvirus) <i>European wheat striate mosaic virus</i> (EWSMV) (Tenuivirus) <i>Euphorbia mosaic virus</i> (EuMV) (Geminiviridae, bigeminivirus) <i>Euphorbia ringspot virus</i> (EuRV) (Potyviridae, potyvirus) <i>Faba bean necrotic yellows virus</i> (FBNYV) (Nanovirus) <i>Festuca leaf streak virus</i> (FLSV) (Rhabdoviridae, cytorhabdovirus) <i>Fig virus S</i> (FVS) (Carlavirus) <i>Figwort mosaic virus</i> (FMV) (Caulimoviridae, caulimovirus) <i>Foxtail mosaic virus</i> (FoMV) (Potexvirus) <i>Frangipani mosaic virus</i> (FrMV) (Tobamovirus) <i>French bean mosaic virus</i> <i>Freesia leaf necrosis virus</i> (FLNV) (Varicosavirus) <i>Freesia mosaic virus</i> (FreMV) (Potyviridae, Potyvirus) <i>Galinsoga mosaic virus</i> (GaMV) (Tombusviridae, carmovirus) <i>Glycine mosaic virus</i> (GMV) (Comoviridae, Comovirus) <i>Grapevine Algerian latent virus</i> (GALV) (Tombusviridae, Tombusvirus) <i>Grapevine asteroid mosaic virus</i> <i>Grapevine berry inner necrosis virus</i> (GINV) (Trichovirus) <i>Grapevine Bulgarian latent virus</i> (GBLV) (Comoviridae, Nepovirus) <i>Grapevine chrome mosaic virus</i> (GCMV) (Comoviridae, Nepovirus) <i>Grapevine leafroll-associated virus 1</i> (GLRAV-1)(Closteroviridae, Ampelovirus) <i>Grapevine stem pitting associated closterovirus</i> (Closteroviridae, Closterovirus) <i>Grapevine Tunisian ringspot virus</i> (GTRSV) (Comoviridae, Nepovirus) <i>Grapevine fanleaf virus</i> (GFLV) (Comoviridae, nepovirus) <i>Grapevine fleck virus</i> (GFkV) <i>Grapevine stunt virus</i> <i>Grapevine vein necrosis virus</i> <i>Grapevine virus A</i> (GVA) (Betaflexiviridae, Vitivirus) <i>Grapevine virus B</i> (GVB) (Vitivirus, Flexiviridae) <i>Grapevine yellow speckle viroid</i> (GYSVd) (Pospiviroidae, Apscaviroid) <i>Groundnut chlorotic leaf streak virus</i> <i>Groundnut chlorotic spotting virus</i> <i>Groundnut rosette assistor virus</i> (GRAV) (Luteoviridae, Enamovirus) <i>Groundnut rosette virus</i> (GRV) (Umbravirus)</p>	<p><i>Datura enation mosaic virus</i> <i>Dendrobium mosaic virus</i> (DeMV) (Potyviridae, potyvirus) <i>Desmodium mosaic virus</i> (DesMV) (Potyviridae, potyvirus) <i>Desmodium yellow mottle virus</i> (DYMV) (Tymovirus) <i>Digitaria streak virus</i> (DSV) (Geminiviridae, monogeminivirus) <i>Digitaria striate mosaic</i> (DiSMV) (Geminiviridae, monogeminivirus) <i>Dioscorea alata virus</i> (Potyviridae, potyvirus) <i>Dioscorea bacilliform virus</i> (DBV) (Caulimoviridae, badnavirus) <i>Dioscorea green banding mosaic virus</i> (Potyviridae, potyvirus) <i>Dodonaea yellows-associated virus</i> <i>Dolichos yellow mosaic virus</i> (DoMYV) (Geminiviridae, bigeminivirus) <i>Dulcamara mottle virus</i> (DuMV) (Tymovirus) <i>Eggplant green mosaic virus</i> (EGMV) (Potyviridae, potyvirus) <i>Eggplant mottled dwarf virus</i> (EMDV) (Rhabdoviridae, nucleorhabdovirus) <i>Elderberry virus</i> (EV) (Carlavirus 1) <i>Epirus cherry virus</i> (EpCV) (Ourmiavirus) <i>Erysimum latent virus</i> (ErLV) (Tymovirus) <i>Echinochloa ragged stunt virus</i> (ERSV) (Reoviridae, Oryzavirus) <i>Elm mottle virus</i> (EMoV) (Bromoviridae, Ilarvirus) <i>European wheat striate mosaic virus</i> (EWSMV) (Tenuivirus) <i>Euphorbia mosaic virus</i> (EuMV) (Geminiviridae, bigeminivirus) <i>Euphorbia ringspot virus</i> (EuRV) (Potyviridae, potyvirus) <i>Faba bean necrotic yellows virus</i> (FBNYV) (Nanovirus) <i>Festuca leaf streak virus</i> (FLSV) (Rhabdoviridae, cytorhabdovirus) <i>Fig virus S</i> (FVS) (Carlavirus) <i>Figwort mosaic virus</i> (FMV) (Caulimoviridae, caulimovirus) <i>Foxtail mosaic virus</i> (FoMV) (Potexvirus) <i>Frangipani mosaic virus</i> (FrMV) (Tobamovirus) <i>French bean mosaic virus</i> <i>Freesia leaf necrosis virus</i> (FLNV) (Varicosavirus) <i>Freesia mosaic virus</i> (FreMV) (Potyviridae, Potyvirus) <i>Galinsoga mosaic virus</i> (GaMV) (Tombusviridae, carmovirus) <i>Glycine mosaic virus</i> (GMV) (Comoviridae, Comovirus) <i>Grapevine Algerian latent virus</i> (GALV) (Tombusviridae, Tombusvirus) <i>Grapevine asteroid mosaic virus</i> <i>Grapevine berry inner necrosis virus</i> (GINV) (Trichovirus) <i>Grapevine Bulgarian latent virus</i> (GBLV) (Comoviridae, Nepovirus) <i>Grapevine chrome mosaic virus</i> (GCMV) (Comoviridae, Nepovirus) <i>Grapevine leafroll-associated virus 1</i> (GLRAV-1)(Closteroviridae, Ampelovirus) <i>Grapevine stem pitting associated closterovirus</i> (Closteroviridae, Closterovirus) <i>Grapevine Tunisian ringspot virus</i> (GTRSV) (Comoviridae, Nepovirus) <i>Grapevine fanleaf virus</i> (GFLV) (Comoviridae, nepovirus) <i>Grapevine fleck virus</i> (GFkV) <i>Grapevine stunt virus</i> <i>Grapevine vein necrosis virus</i> <i>Grapevine virus A</i> (GVA) (Betaflexiviridae, Vitivirus) <i>Grapevine virus B</i> (GVB) (Vitivirus, Flexiviridae) <i>Grapevine yellow speckle viroid</i> (GYSVd) (Pospiviroidae, Apscaviroid) <i>Groundnut chlorotic leaf streak virus</i> <i>Groundnut chlorotic spotting virus</i> <i>Groundnut rosette assistor virus</i> (GRAV) (Luteoviridae, Enamovirus) <i>Groundnut rosette virus</i> (GRV) (Umbravirus)</p>	
--	--	--

<p>Groundnut eyespot virus (GEV) (Potyviridae, Potyvirus) Guar symptomless virus (GSLV) (Potyviridae, Potyvirus) Guar top necrosis virus Guinea grass mosaic virus (GGMV) (Potyviridae, Potyvirus) Helenium virus S (HVS) (Carlavirus) Helenium virus Y (HVY) (Potyviridae, Potyvirus) Henbane mosaic virus (HBMV) (Potyviridae, Potyvirus) Heracleum latent virus (HLV) (Vitivirus) Hibiscus latent ringspot virus (HLRSV) (Comoviridae, nepovirus) Hibiscus chlorotic ringspot virus (HCRSV) (Tombusviridae, Carmovirus) Hippeastrum mosaic virus Honeysuckle latent virus (LnLV) (Carlavirus) Hop American latent (AHLV) (Carlavirus) Hop latent virus (HpLV) (Carlavirus) Hop mosaic virus (HpMV) (Carlavirus) Hop trefoil cryptic virus 1 (HTCV-1) (Partitiviridae, alphacryptovirus) Hop trefoil cryptic virus 2 (HTCV-2) (Partitiviridae, betacryptovirus) Hop trefoil cryptic virus 3 (HTCV-3) (Partitiviridae, alphacryptovirus) Hop stunt viroid (HpSVd) (Hostuviroid, Pospiviroidae) Hordeum mosaic virus (HoMV) (Potyviridae, Potyvirus) Humulus japonicus virus (HJV) (Bromoviridae, ilarvirus) Hyacinth mosaic virus (HyaMV) (Potyviridae, Potyvirus) Hydrangea mosaic virus (HdMV) (Bromoviridae, ilarvirus) Hydrangea ringspot virus (HRSV) (Potexvirus) Horsegram yellow mosaic virus (HgYMV) (Geminiviridae, Begomovirus) Iris fulva mosaic virus (IFMV) (Potyviridae, Potyvirus) Iris Japanese necrotic ring virus Iris mild mosaic virus (IMMV) (Potyviridae, Potyvirus) Iris severe mosaic virus (ISMV) (Potyviridae, Potyvirus) Impatiens necrotic spot virus (INSV) (Bunyaviridae, Tospovirus) Indian peanut clump virus (IPCV) (Pecluvirus) Indonesian soybean dwarf virus (ISDV) (Luteoviridae, Enamovirus) Iranian maize mosaic virus Jatropha mosaic virus (JMV) (Geminiviridae, bigeminivirus) Johnsongrass mosaic (JGMV) (Potyviridae, Potyvirus) Kalanchoe isometric virus Kalanchoe latent virus (KLV) (Carlavirus) Kalanchoe top-spotting virus (KTSV) (Caulimoviridae, badnavirus) Kennedy yellow mosaic virus (KYMV) (Tymovirus) Kennedy virus Y (KVY) (Potyviridae, Potyvirus) Kyuri green mottle mosaic virus (KGMMV) (Tobamovirus) Konjac mosaic virus (KoMV) (Potyviridae, Potyvirus) Kober stem grooving Lamium mild mottle virus (LMMV) (Comoviridae, fabavirus) Lato River virus (LRV) (Tombusviridae, tombusvirus) Leek yellow stripe virus (LYSV) (Potyviridae, potyvirus) Lettuce big-vein virus (LBVV) (Varicosavirus) Lettuce mosaic virus (LMV) (Potyviridae, potyvirus) Lettuce necrotic yellows virus (LNYV) (Rhabdoviridae, cytorhabdovirus) Lettuce speckles mottle virus (LSMV) (Umbravirus) Lilac chlorotic leafspot virus (LCLV) (Capillovirus) Lilac mottle virus (LiMV) (Carlavirus)</p>	<p>Groundnut eyespot virus (GEV) (Potyviridae, Potyvirus) Guar symptomless virus (GSLV) (Potyviridae, Potyvirus) Guar top necrosis virus Guinea grass mosaic virus (GGMV) (Potyviridae, Potyvirus) Helenium virus S (HVS) (Carlavirus) Helenium virus Y (HVY) (Potyviridae, Potyvirus) Henbane mosaic virus (HBMV) (Potyviridae, Potyvirus) Heracleum latent virus (HLV) (Vitivirus) Hibiscus latent ringspot virus (HLRSV) (Comoviridae, nepovirus) Hibiscus chlorotic ringspot virus (HCRSV) (Tombusviridae, Carmovirus) Hippeastrum mosaic virus Honeysuckle latent virus (LnLV) (Carlavirus) Hop American latent (AHLV) (Carlavirus) Hop latent virus (HpLV) (Carlavirus) Hop mosaic virus (HpMV) (Carlavirus) Hop trefoil cryptic virus 1 (HTCV-1) (Partitiviridae, alphacryptovirus) Hop trefoil cryptic virus 2 (HTCV-2) (Partitiviridae, betacryptovirus) Hop trefoil cryptic virus 3 (HTCV-3) (Partitiviridae, alphacryptovirus) Hop stunt viroid (HpSVd) (Hostuviroid, Pospiviroidae) Hordeum mosaic virus (HoMV) (Potyviridae, Potyvirus) Humulus japonicus virus (HJV) (Bromoviridae, ilarvirus) Hyacinth mosaic virus (HyaMV) (Potyviridae, Potyvirus) Hydrangea mosaic virus (HdMV) (Bromoviridae, ilarvirus) Hydrangea ringspot virus (HRSV) (Potexvirus) Horsegram yellow mosaic virus (HgYMV) (Geminiviridae, Begomovirus) Iris fulva mosaic virus (IFMV) (Potyviridae, Potyvirus) Iris Japanese necrotic ring virus Iris mild mosaic virus (IMMV) (Potyviridae, Potyvirus) Iris severe mosaic virus (ISMV) (Potyviridae, Potyvirus) Impatiens necrotic spot virus (INSV) (Bunyaviridae, Tospovirus) Indian peanut clump virus (IPCV) (Pecluvirus) Indonesian soybean dwarf virus (ISDV) (Luteoviridae, Enamovirus) Iranian maize mosaic virus Jatropha mosaic virus (JMV) (Geminiviridae, bigeminivirus) Johnsongrass mosaic (JGMV) (Potyviridae, Potyvirus) Kalanchoe isometric virus Kalanchoe latent virus (KLV) (Carlavirus) Kalanchoe top-spotting virus (KTSV) (Caulimoviridae, badnavirus) Kennedy yellow mosaic virus (KYMV) (Tymovirus) Kennedy virus Y (KVY) (Potyviridae, Potyvirus) Kyuri green mottle mosaic virus (KGMMV) (Tobamovirus) Konjac mosaic virus (KoMV) (Potyviridae, Potyvirus) Kober stem grooving Lamium mild mottle virus (LMMV) (Comoviridae, fabavirus) Lato River virus (LRV) (Tombusviridae, tombusvirus) Leek yellow stripe virus (LYSV) (Potyviridae, potyvirus) Lettuce big-vein virus (LBVV) (Varicosavirus) Lettuce mosaic virus (LMV) (Potyviridae, potyvirus) Lettuce necrotic yellows virus (LNYV) (Rhabdoviridae, cytorhabdovirus) Lettuce speckles mottle virus (LSMV) (Umbravirus) Lilac chlorotic leafspot virus (LCLV) (Capillovirus) Lilac mottle virus (LiMV) (Carlavirus)</p>	
---	---	--

<p><i>Lilac ring mottle virus (LRMV) (Bromoviridae, ilarvirus)</i> <i>Lily mottle virus (LMOV) (Potyviridae, potyvirus)</i> <i>Lily virus X (LVX) (Potexvirus)</i> <i>Lettuce infectious yellows virus (LIYV) (Closteroviridae, Crinivirus)</i> <i>Limabean golden mosaic virus (LGMV) (Geminiviridae, Begomovirus)</i> <i>Little cherry virus (LChV) (Closteroviridae, Closterovirus)</i> <i>LN33 stem grooving</i> <i>Lucerne Australian symptomless virus (LASV) (Comoviridae, Nepovirus)</i> <i>Lucerne vein yellowing virus</i> <i>Lucerne transient streak virus (LTSV) (Sobemovirus)</i> <i>Lychnis ringspot virus (LRSV) (Hordeivirus)</i> <i>Maclura mosaic virus (MacMV) (Potyviridae, macluravirus)</i> <i>Maize chlorotic dwarf virus (MCDV) (Sequiviridae, waikavirus)</i> <i>Maize chlorotic mottle virus (MCMV) (Tombusviridae, machlomovirus)</i> <i>Maize dwarf mosaic virus (MDMV) (Potyviridae, potyvirus)</i> <i>Maize line virus</i> <i>Maize mosaic virus (MMV) (Rhabdoviridae, nucleorhabdovirus)</i> <i>Maize rayado fino virus (MRFV) (Marafivirus)</i> <i>Maize white line mosaic virus (MWLMV)</i> <i>Maize mottle/chlorotic stunt virus</i> <i>Maize rough dwarf virus (MRDV) (Reoviridae, Fijivirus)</i> <i>Maize streak virus (MSV) (Genimiviridae, Mastrevirus)</i> <i>Malvastrum mottle virus</i> <i>Malva vein clearing virus (MVCV) (Potyviridae, potyvirus)</i> <i>Marigold mottle virus (MaMoV) (Potyviridae, potyvirus)</i> <i>Melandrium yellow fleck virus (MYFV) (Bromoviridae, bromovirus)</i> <i>Melon leaf curl virus (MLCV) (Geminiviridae, bigeminivirus)</i> <i>Melon necrotic spot virus (MNSV) (Tombusviridae, carmovirus)</i> <i>Melon Ourmia virus (OuMV) (Ourmiavirus)</i> <i>Melon rugose mosaic virus (MRMV) (Tymovirus)</i> <i>Milk vetch dwarf virus (MDV) (Nanovirus)</i> <i>Mimosa mosaic virus</i> <i>Mirabilis mosaic virus (MiMV) (Caulimovirus)</i> <i>Miscanthus streak virus (MiSV) (Geminiviridae, monogeminivirus)</i> <i>Mulberry latent virus (MLV) (Carlavirus)</i> <i>Mulberry ringspot virus (MRSV) (Comoviridae, nepovirus)</i> <i>Mung bean yellow mosaic virus (MYMV) (Geminiviridae, bigeminivirus)</i> <i>Muskmelon vein necrosis virus (MuVNV) (Carlavirus)</i> <i>Myrobalan latent ringspot virus (MLRSV) (Comoviridae, nepovirus)</i> <i>Narcissus latent virus (NLV) (Potyviridae, macluravirus)</i> <i>Narcissus mosaic virus (NMV) (Potexvirus)</i> <i>Narcissus yellow stripe virus (Potyviridae, potyvirus)</i> <i>Natsudaidai dwarf virus (NDV) (Sadwavirus)</i> <i>Neckar River virus (NRV) (Tombusviridae, tombusvirus)</i> <i>Nerine latent virus (NeLV) (Carlavirus)</i> <i>Nerine virus (NV) (Potyviridae, potyvirus)</i> <i>Nerine virus X (NVX) (Potexvirus)</i> <i>Northern cereal mosaic virus (NCMV) (Rhabdoviridae, Cytorhabdovirus)</i> <i>Oak ringspot virus</i> <i>Oat blue dwarf virus (OBDV) (Marafivirus)</i> <i>Oat golden stripe virus (OGSV) (Furovirus)</i> <i>Oat mosaic virus (OMV)(Potyviridae, bymovirus)</i></p>	<p><i>Lilac ring mottle virus (LRMV) (Bromoviridae, ilarvirus)</i> <i>Lily mottle virus (LMOV) (Potyviridae, potyvirus)</i> <i>Lily virus X (LVX) (Potexvirus)</i> <i>Lettuce infectious yellows virus (LIYV) (Closteroviridae, Crinivirus)</i> <i>Limabean golden mosaic virus (LGMV) (Geminiviridae, Begomovirus)</i> <i>Little cherry virus (LChV) (Closteroviridae, Closterovirus)</i> <i>LN33 stem grooving</i> <i>Lucerne Australian symptomless virus (LASV) (Comoviridae, Nepovirus)</i> <i>Lucerne vein yellowing virus</i> <i>Lucerne transient streak virus (LTSV) (Sobemovirus)</i> <i>Lychnis ringspot virus (LRSV) (Hordeivirus)</i> <i>Maclura mosaic virus (MacMV) (Potyviridae, macluravirus)</i> <i>Maize chlorotic dwarf virus (MCDV) (Sequiviridae, waikavirus)</i> <i>Maize chlorotic mottle virus (MCMV) (Tombusviridae, machlomovirus)</i> <i>Maize dwarf mosaic virus (MDMV) (Potyviridae, potyvirus)</i> <i>Maize line virus</i> <i>Maize mosaic virus (MMV) (Rhabdoviridae, nucleorhabdovirus)</i> <i>Maize rayado fino virus (MRFV) (Marafivirus)</i> <i>Maize white line mosaic virus (MWLMV)</i> <i>Maize mottle/chlorotic stunt virus</i> <i>Maize rough dwarf virus (MRDV) (Reoviridae, Fijivirus)</i> <i>Maize streak virus (MSV) (Genimiviridae, Mastrevirus)</i> <i>Malvastrum mottle virus</i> <i>Malva vein clearing virus (MVCV) (Potyviridae, potyvirus)</i> <i>Marigold mottle virus (MaMoV) (Potyviridae, potyvirus)</i> <i>Melandrium yellow fleck virus (MYFV) (Bromoviridae, bromovirus)</i> <i>Melon leaf curl virus (MLCV) (Geminiviridae, bigeminivirus)</i> <i>Melon necrotic spot virus (MNSV) (Tombusviridae, carmovirus)</i> <i>Melon Ourmia virus (OuMV) (Ourmiavirus)</i> <i>Melon rugose mosaic virus (MRMV) (Tymovirus)</i> <i>Milk vetch dwarf virus (MDV) (Nanovirus)</i> <i>Mimosa mosaic virus</i> <i>Mirabilis mosaic virus (MiMV) (Caulimovirus)</i> <i>Miscanthus streak virus (MiSV) (Geminiviridae, monogeminivirus)</i> <i>Mulberry latent virus (MLV) (Carlavirus)</i> <i>Mulberry ringspot virus (MRSV) (Comoviridae, nepovirus)</i> <i>Mung bean yellow mosaic virus (MYMV) (Geminiviridae, bigeminivirus)</i> <i>Muskmelon vein necrosis virus (MuVNV) (Carlavirus)</i> <i>Myrobalan latent ringspot virus (MLRSV) (Comoviridae, nepovirus)</i> <i>Narcissus latent virus (NLV) (Potyviridae, macluravirus)</i> <i>Narcissus mosaic virus (NMV) (Potexvirus)</i> <i>Narcissus yellow stripe virus (Potyviridae, potyvirus)</i> <i>Natsudaidai dwarf virus (NDV) (Sadwavirus)</i> <i>Neckar River virus (NRV) (Tombusviridae, tombusvirus)</i> <i>Nerine latent virus (NeLV) (Carlavirus)</i> <i>Nerine virus (NV) (Potyviridae, potyvirus)</i> <i>Nerine virus X (NVX) (Potexvirus)</i> <i>Northern cereal mosaic virus (NCMV) (Rhabdoviridae, Cytorhabdovirus)</i> <i>Oak ringspot virus</i> <i>Oat blue dwarf virus (OBDV) (Marafivirus)</i> <i>Oat golden stripe virus (OGSV) (Furovirus)</i> <i>Oat mosaic virus (OMV)(Potyviridae, bymovirus)</i></p>	
---	---	--

<p>Oat necrotic mottle virus (ONMV) (Potyviridae, rymovirus) Oat red streak mosaic virus Oat sterile dwarf virus (OSDV) (Reoviridae, Fijivirus) Okra leaf-curl virus (OLCV) (Geminivirus, bigeminivirus) Okra mosaic virus (OkMV) (Tymovirus) Olive latent ringspot virus (OLRSV) (Comoviridae, nepovirus) Ononis yellow mosaic virus (OYMV) (Tymovirus) Opuntia Sammons' virus (SOV) (Tobamovirus) Ornithogalum mosaic virus Pangola stunt virus (PaSV) (Reoviridae, Fijivirus) Panicum streak virus (PanSV) (Geminiviridae, monogeminivirus) Papaya leaf curl virus (PaLCuV) (Begomovirus, Geminiviridae) Papaya meleira virus (PMeV) Papaya mosaic virus (PapMV) (Potexvirus, Flexiviridae) Paprika mild mottle virus (PaMMV) (Tobamovirus) Parietaria mottle virus (PmoV) (Bromoviridae, ilarvirus) Parsnip leafcurl virus Parsnip mosaic virus (ParMV) Parsnip yellow fleck virus (PYFV) (Sequiviridae, sequivirus) Paspalum striate mosaic virus (PSMV) (Geminiviridae, monogeminivirus) Passiflora latent virus (PLV)(Carlavirus) Passiflora South African virus (Potyviridae, potyvirus) Passiflora ringspot virus (PFRSV) (Potyviridae, potyvirus) Passionfruit woodiness virus (PWV) (Potyviridae, potyvirus) Passionfruit yellow mosaic virus (PaYMV) (Tymovirus) Patchouli mosaic virus (Potyviridae, potyvirus) Pea early browning virus (PEBV) (Tobravirus) Pea enation mosaic virus-1 (PEMV-1) (Luteovirus) Pea enation mosaic virus-2 (PEMV-2) (Umbravirus) Pea mild mosaic virus (PmiMV) (Comoviridae, comovirus) Pea mosaic virus (Potyviridae, potyvirus) Peanut chlorotic streak virus (PCSV) (Caulimoviridae, caulimovirus) Peanut green mosaic virus (Potyviridae, potyvirus) Peanut stunt virus (PSV) (Bromoviridae, cucumovirus) Peanut yellow spot virus (PeYMV) (Tospovirus) Peach American mosaic virus Peach latent mosaic viroid (PLMVd) Peach mosaic virus (PMV) (Trichovirus) Peach rosette mosaic virus (PRMV) (Comoviridae, Nepovirus) Peanut clump virus (PCV) (Pecluvirus) Pear blister canker viroid (PBCVd) (Pospiviroidae, Apscaviriod) Pear necrotic spot virus</p> <p>Pineapple bacilliform virus (PBV) (Badnavirus, Caulimoviridae) Pineapple mealybug wilt-associated virus-1 (PMWaV-1) (Ampelovirus, Closteroviridae) Pineapple mealybug wilt-associated virus-2 (PMWaV-2) (Ampelovirus, Closteroviridae) Plum American line pattern virus (APLPV) (Ilarvirus, Bromoviridae) Plum bark split virus Plum pox virus (PPV) (Potyviridae, Potyvirus) Plum rusty blotch</p>	<p>Oat necrotic mottle virus (ONMV) (Potyviridae, rymovirus) Oat red streak mosaic virus Oat sterile dwarf virus (OSDV) (Reoviridae, Fijivirus) Okra leaf-curl virus (OLCV) (Geminivirus, bigeminivirus) Okra mosaic virus (OkMV) (Tymovirus) Olive latent ringspot virus (OLRSV) (Comoviridae, nepovirus) Ononis yellow mosaic virus (OYMV) (Tymovirus) Opuntia Sammons' virus (SOV) (Tobamovirus) Ornithogalum mosaic virus Pangola stunt virus (PaSV) (Reoviridae, Fijivirus) Panicum streak virus (PanSV) (Geminiviridae, monogeminivirus) Papaya leaf curl virus (PaLCuV) (Begomovirus, Geminiviridae) Papaya meleira virus (PMeV) Papaya mosaic virus (PapMV) (Potexvirus, Flexiviridae) Paprika mild mottle virus (PaMMV) (Tobamovirus) Parietaria mottle virus (PmoV) (Bromoviridae, ilarvirus) Parsnip leafcurl virus Parsnip mosaic virus (ParMV) Parsnip yellow fleck virus (PYFV) (Sequiviridae, sequivirus) Paspalum striate mosaic virus (PSMV) (Geminiviridae, monogeminivirus) Passiflora latent virus (PLV)(Carlavirus) Passiflora South African virus (Potyviridae, potyvirus) Passiflora ringspot virus (PFRSV) (Potyviridae, potyvirus) Passionfruit woodiness virus (PWV) (Potyviridae, potyvirus) Passionfruit yellow mosaic virus (PaYMV) (Tymovirus) Patchouli mosaic virus (Potyviridae, potyvirus) Pea early browning virus (PEBV) (Tobravirus) Pea enation mosaic virus-1 (PEMV-1) (Luteovirus) Pea enation mosaic virus-2 (PEMV-2) (Umbravirus) Pea mild mosaic virus (PmiMV) (Comoviridae, comovirus) Pea mosaic virus (Potyviridae, potyvirus) Peanut chlorotic streak virus (PCSV) (Caulimoviridae, caulimovirus) Peanut green mosaic virus (Potyviridae, potyvirus) Peanut stunt virus (PSV) (Bromoviridae, cucumovirus) Peanut yellow spot virus (PeYMV) (Tospovirus) Peach American mosaic virus Peach latent mosaic viroid (PLMVd) Peach mosaic virus (PMV) (Trichovirus) Peach rosette mosaic virus (PRMV) (Comoviridae, Nepovirus) Peanut clump virus (PCV) (Pecluvirus) Pear blister canker viroid (PBCVd) (Pospiviroidae, Apscaviriod) Pear necrotic spot virus <u>Pepper chat fruit viroid (PCFVd) (Pospiviroidae, Pospiviroid)</u> Pineapple bacilliform virus (PBV) (Badnavirus, Caulimoviridae) Pineapple mealybug wilt-associated virus-1 (PMWaV-1) (Ampelovirus, Closteroviridae) Pineapple mealybug wilt-associated virus-2 (PMWaV-2) (Ampelovirus, Closteroviridae) Plum American line pattern virus (APLPV) (Ilarvirus, Bromoviridae) Plum bark split virus Plum pox virus (PPV) (Potyviridae, Potyvirus) Plum rusty blotch</p>	
---	---	--

<p>Potato black ringspot virus (PBRSV) (Comoviridae, Nepovirus) Potato deforming mosaic virus Potato mop-top virus (PMTV) (Pomovirus) Potato virus T (PVT) (Trichovirus) Potato virus U (PVU) (Comoviridae, Nepovirus) Potato virus V (PVV) (Potyviridae, Potyvirus) Potato yellow dwarf virus (PYDV) (Rhabdoviridae, Nucleorhabdovirus) Potato yellow vein virus (PYVV) (Closteroviridae, Crinivirus) Potato yellowing virus Prune dwarf virus (PDV) (Bromoviridae, Ilarvirus) Prunus necrotic ringspot virus (PNRSV) (Bromoviridae, Ilarvirus) Raspberry leaf curl virus Raspberry ringspot virus (RpRSV) (Comoviridae, Nepovirus) Red clover mottle virus (RCMV) (Comoviridae, Comovirus) Rice gall dwarf virus (RGDV) (Reoviridae, Phytoreovirus) Rice tungro bacilliform virus (RTBV) (Caulimoviridae, Rice tungro bacilliform-like virus) Rice tungro spherical virus (RTSV) (Sequiviridae, Waikavirus) Rice yellow mottle virus (RYMV) (Sobemovirus) Rupestris stem pitting-associated virus (Foveavirus, Flexiviridae) Squash leaf curl virus (SLCV) (Geminiviridae, Begomovirus) Squash mosaic virus (SqMV) (Comoviridae, Comovirus) Strawberry latent ringspot virus (SLRSV) (Sadwavirus) Strawberry veinbanding virus (SVBV) (Caulimoviridae, Caulimovirus) Sugarcane yellow leaf virus (ScYLV) (Polerovirus, Luteoviridae) Tobacco rattle virus (TRV) (Tobravirus) Tobacco ringspot virus (TRSV) (Comoviridae, Nepovirus) Tobacco mosaic virus (TMV) (Tobamovirus) Tobacco necrosis virus Tobacco streak virus (TSV) (Ilarvirus, Bromoviridae)</p> <p>Tomato black ring virus (TBRV) (Comoviridae, Nepovirus) Tomato bushy stunt virus (TBSV) (Tombusvirus, Tombusviridae)</p> <p>Tomato mottle virus (ToMoV) (Geminiviridae, Begomovirus)</p> <p>Tomato ringspot virus (ToRSV) (Comoviridae, Nepovirus)</p> <p>Tulare apple mosaic virus (TAMV) (Ilarvirus, Bromoviridae) Tulip breaking virus Ullucus virus C (UVC) (Comoviridae, comovirus) Ullucus mild mottle virus (UMMV) (Tobamovirus) Ullucus mosaic virus (UMV) (Potyviridae, potyvirus) Urd bean leaf crinkle virus Vallota mosaic virus (ValMV) (Potyviridae, potyvirus) Vanilla mosaic virus (VanMV) (Potyviridae, potyvirus) Vanilla necrosis virus (Potyviridae, potyvirus) Velvet tobacco mottle virus (VTMoV) (Sobemovirus) Viola mottle virus (VMV) (Potexvirus) Voandzeia distortion mosaic virus (Potyviridae, potyvirus) Voandzeia necrotic mosaic virus (VNMV) (Tymovirus)</p>	<p>Potato black ringspot virus (PBRSV) (Comoviridae, Nepovirus) Potato deforming mosaic virus Potato mop-top virus (PMTV) (Pomovirus) Potato virus T (PVT) (Trichovirus) Potato virus U (PVU) (Comoviridae, Nepovirus) Potato virus V (PVV) (Potyviridae, Potyvirus) Potato yellow dwarf virus (PYDV) (Rhabdoviridae, Nucleorhabdovirus) Potato yellow vein virus (PYVV) (Closteroviridae, Crinivirus) Potato yellowing virus Prune dwarf virus (PDV) (Bromoviridae, Ilarvirus) Prunus necrotic ringspot virus (PNRSV) (Bromoviridae, Ilarvirus) Raspberry leaf curl virus Raspberry ringspot virus (RpRSV) (Comoviridae, Nepovirus) Red clover mottle virus (RCMV) (Comoviridae, Comovirus) Rice gall dwarf virus (RGDV) (Reoviridae, Phytoreovirus) Rice tungro bacilliform virus (RTBV) (Caulimoviridae, Rice tungro bacilliform-like virus) Rice tungro spherical virus (RTSV) (Sequiviridae, Waikavirus) Rice yellow mottle virus (RYMV) (Sobemovirus) Rupestris stem pitting-associated virus (Foveavirus, Flexiviridae) Squash leaf curl virus (SLCV) (Geminiviridae, Begomovirus) Squash mosaic virus (SqMV) (Comoviridae, Comovirus) Strawberry latent ringspot virus (SLRSV) (Sadwavirus) Strawberry veinbanding virus (SVBV) (Caulimoviridae, Caulimovirus) Sugarcane yellow leaf virus (ScYLV) (Polerovirus, Luteoviridae) Tobacco rattle virus (TRV) (Tobravirus) Tobacco ringspot virus (TRSV) (Comoviridae, Nepovirus) Tobacco mosaic virus (TMV) (Tobamovirus) Tobacco necrosis virus Tobacco streak virus (TSV) (Ilarvirus, Bromoviridae) <u>Tomato apical stunt viroid (TASVd) (Pospiviroidae, Pospiviroid)</u> Tomato black ring virus (TBRV) (Comoviridae, Nepovirus) Tomato bushy stunt virus (TBSV) (Tombusvirus, Tombusviridae) <u>Tomato chlorotic dwarf viroid (TCDVd) (Pospiviroidae, Pospiviroid)</u> <u>Tomato mottle mosaic virus (ToMMV) (Virgaviridae, Tobamovirus)</u> Tomato mottle virus (ToMoV) (Geminiviridae, Begomovirus) <u>Tomato planta macho viroid (TPMVd) (Pospiviroidae, Pospiviroid)</u> Tomato ringspot virus (ToRSV) (Comoviridae, Nepovirus) <u>Tomato yellow leaf curl virus (TYLCV) (Geminiviridae, Begomovirus)</u> Tulare apple mosaic virus (TAMV) (Ilarvirus, Bromoviridae) Tulip breaking virus Ullucus virus C (UVC) (Comoviridae, comovirus) Ullucus mild mottle virus (UMMV) (Tobamovirus) Ullucus mosaic virus (UMV) (Potyviridae, potyvirus) Urd bean leaf crinkle virus Vallota mosaic virus (ValMV) (Potyviridae, potyvirus) Vanilla mosaic virus (VanMV) (Potyviridae, potyvirus) Vanilla necrosis virus (Potyviridae, potyvirus) Velvet tobacco mottle virus (VTMoV) (Sobemovirus) Viola mottle virus (VMV) (Potexvirus) Voandzeia distortion mosaic virus (Potyviridae, potyvirus) Voandzeia necrotic mosaic virus (VNMV) (Tymovirus)</p>	
---	--	--

<p>Watercress yellow spot virus (WYSV) Watermelon chlorotic stunt virus (WmCSV) (Geminiviridae, bigeminivirus) Watermelon curly mottle virus (WmCMV) (Geminiviridae, bigeminivirus) Watermelon mosaic virus 2 (WMV-2) (Potyviridae, potyvirus) Wheat American striate mosaic virus (WASMV) (Rhabdovirus, ucleorhabdovirus) Wheat yellow leaf virus (WYLV) (Closteroviridae, Closterovirus) Wheat dwarf virus (WDV) (Geminiviridae, monogeminivirus) Wheat soil-borne mosaic virus (SBWMV) (Furovirus) Wheat streak mosaic virus (WSMV) (Potyviridae, rymovirus) Wheat spindle streak mosaic virus (WSSMV) (Potyvirida, bymovirus) Wheat yellow mosaic virus (WYMV) (Potyviridae, bymovirus) White clover mosaic virus (WCIMV) (Potexvirus) Wild cucumber mosaic virus (WCMV) (Tymovirus) Wild potato mosaic (WPMV) (Potyviridae, Potyvirus) Wineberry latent virus (WLV) Wisteria vein mosaic virus (WVMV) (Potyviridae, Potyvirus) Yam mosaic virus (YMV) (Potyviridae, Potyvirus) Zucchini yellow fleck virus (ZYFV) (Potyviridae, Potyvirus) 備註：以上病毒之媒介有害生物亦一併列入管制</p>	<p>Watercress yellow spot virus (WYSV) Watermelon chlorotic stunt virus (WmCSV) (Geminiviridae, bigeminivirus) Watermelon curly mottle virus (WmCMV) (Geminiviridae, bigeminivirus) Watermelon mosaic virus 2 (WMV-2) (Potyviridae, potyvirus) Wheat American striate mosaic virus (WASMV) (Rhabdovirus, ucleorhabdovirus) Wheat yellow leaf virus (WYLV) (Closteroviridae, Closterovirus) Wheat dwarf virus (WDV) (Geminiviridae, monogeminivirus) Wheat soil-borne mosaic virus (SBWMV) (Furovirus) Wheat streak mosaic virus (WSMV) (Potyviridae, rymovirus) Wheat spindle streak mosaic virus (WSSMV) (Potyvirida, bymovirus) Wheat yellow mosaic virus (WYMV) (Potyviridae, bymovirus) White clover mosaic virus (WCIMV) (Potexvirus) Wild cucumber mosaic virus (WCMV) (Tymovirus) Wild potato mosaic (WPMV) (Potyviridae, Potyvirus) Wineberry latent virus (WLV) Wisteria vein mosaic virus (WVMV) (Potyviridae, Potyvirus) Yam mosaic virus (YMV) (Potyviridae, Potyvirus) Zucchini yellow fleck virus (ZYFV) (Potyviridae, Potyvirus) 備註：以上病毒之媒介有害生物亦一併列入管制</p>	
<p>十六、(刪除)</p>	<p>十六、自各國輸入茄屬及番椒屬種子應依「茄屬及番椒屬種子輸入檢疫條件」(如附件)辦理。</p>	<p>一、本點刪除。 二、本點所規範之九種病原業經風險評估，分別列於第一點增訂相關檢疫條件，爰刪除本點規定。</p>
<p>十七、本檢疫條件所稱繁殖用生植株，指供種植、繁殖、嫁接、授粉之生植株之全部及部分，包含根、莖、葉、枝條、種球、塊莖、鱗莖、球莖、根狀莖、種子及花粉等。</p>		<p>一、本點新增。 二、為明確本檢疫規定所稱「繁殖用生植株」之定義，爰參考國際植物防疫檢疫措施標準第5號(ISPM No.5)之生植株(plant)及繁殖用生植株(plants for planting)之定義予以增列。</p>

中華民國輸入植物或植物產品檢疫規定乙、有條件輸入植物或植物產品之檢疫條件第十六點附件茄屬及番椒屬種子輸入檢疫條件修正對照表

修正規定	現行規定	說明
	<p>一、輸入茄屬(<i>Solanum</i> spp.)及番椒屬(<i>Capsicum</i> spp.)種子，除依植物防疫檢疫法與其施行細則及中華民國輸入植物或植物產品檢疫規定辦理外，依本檢疫條件辦理。</p>	<p>一、<u>本點刪除</u>。 二、本附件所規範之九種病原業經風險評估，分別列於第一點增訂相關檢疫條件，爰刪除本點規定。</p>
	<p>二、輸入茄屬及番椒屬種子應檢附輸出國植物檢疫機關簽發之植物檢疫證明書，證明於輸出前經實驗室檢測未罹染下列病毒及類病毒：</p> <p>(一) 病毒：番茄斑駁嵌紋病毒 (<i>Tomato mottle mosaic virus</i>, ToMMV)、香瓜茄嵌紋病毒 (<i>Pepino mosaic virus</i>, PepMV) 及番茄黃化捲葉病毒 (<i>Tomato yellow leaf curl virus</i>, TYLCV)。</p> <p>(二) 類病毒：鯨魚藤潛隱類病毒 (<i>Columnea latent viroid</i>, CLVd)、馬鈴薯紡錘型塊莖類病毒 (<i>Potato spindle tuber viroid</i>, PSTVd)、番茄莖頂矮化類病毒 (<i>Tomato apical stunt viroid</i>, TASVd)、番茄褪綠矮化類病毒 (<i>Tomato chlorotic dwarf viroid</i>, TCDVd)、番茄植株雄化類病毒 (<i>Tomato planta</i></p>	<p>一、<u>本點刪除</u>。 二、本附件所規範之九種病原業經風險評，分別列於第一點增訂相關檢疫條件，爰刪除本點規定。</p>

	<p><i>macho viroid</i>, TPMVd) 及 番 椒 小 果 類 病 毒 (<i>Pepper chat fruit viroid</i>, PCFVd)。</p>	
	<p>三、輸入檢疫之程序、方法及抽驗數量，依我國植物防疫檢疫法與其施行細則及相關規定辦理。</p>	<p>一、本點刪除。 二、本附件所規範之九種病原業經風險評估，分別列於第一點增訂相關檢疫條件，爰刪除本點規定。</p>
	<p>四、輸入茄屬及番椒屬種子有下列情形之一者，不得輸入：</p> <p>(一) 未檢附輸出國植物檢疫證明書。</p> <p>(二) 檢附之輸出國植物檢疫證明書證明事項不符第二點規定者，且經植物檢疫機關通知限期補正，屆期未補正或放棄補正。</p>	<p>一、本點刪除。 二、本附件所規範之九種病原業經風險評估，分別列於第一點增訂相關檢疫條件，爰刪除本點規定。</p>